

BinCARD Polyclonal Antibody

Catalog No: YT0491

Reactivity: Human; Mouse

Applications: IHC;IF;ELISA

Target: BinCARD

Gene Name: C9orf89

Protein Name: Bcl10-interacting CARD protein

Q96LW7

Q9D1I2

Human Gene Id: 84270

Human Swiss Prot

No:

Mouse Gene ld: 68480

Mouse Swiss Prot

No:

Immunogen: The antiserum was produced against synthesized peptide derived from human

C9orf89. AA range:21-70

Specificity: BinCARD Polyclonal Antibody detects endogenous levels of BinCARD protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution : IHC 1:100 - 1:300. ELISA: 1:40000.. IF 1:50-200

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)



Molecularweight:

26kD

Background:

function:Plays a role in inhibiting the effects of BCL10-induced activation of NF-kappa-B. May inhibit the phosphorylation of BCL10 in a CARD-dependent manner.,PTM:Isoform 2 is phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 CARD domain.,subcellular location:Co-expression with BCL10 induced translocation from nucleus to cytosol.,subunit:Associates with BCL10 by CARD-CARD interaction.,tissue specificity:Expressed in ovary, testis, placenta, skeletal muscle, kidney, lung, heart and liver (at protein level). Expressed in thymus and brain.,

Function:

function:Plays a role in inhibiting the effects of BCL10-induced activation of NF-kappa-B. May inhibit the phosphorylation of BCL10 in a CARD-dependent manner.,PTM:Isoform 2 is phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 CARD domain.,subcellular location:Co-expression with BCL10 induced translocation from nucleus to cytosol.,subunit:Associates with BCL10 by CARD-CARD interaction.,tissue specificity:Expressed in ovary, testis, placenta, skeletal muscle, kidney, lung, heart and liver (at protein level). Expressed in thymus and brain.,

Subcellular Location:

[Isoform 1]: Nucleus . Coexpression with BCL10 induced translocation from nucleus to cytosol.; [Isoform 2]: Endoplasmic reticulum membrane ; Single-pass membrane protein . Mitochondrion membrane ; Single-pass membrane protein .

Expression:

Expressed in ovary, testis, placenta, skeletal muscle, kidney, lung, heart and liver (at protein level). Expressed in thymus and brain.


Sort:

2699

No4:

1

Products Images



Immunohistochemistry analysis of paraffin-embedded human thyroid gland tissue, using C9orf89 Antibody. The picture on the right is blocked with the synthesized peptide.