

## HBP1 Polyclonal Antibody


Catalog No :	YT2103
Reactivity :	Human;Mouse;Rat;Monkey
Applications :	WB;IHC;IF;ELISA
Target :	HBP1
Gene Name :	HBP1
Protein Name :	HMG box-containing protein 1
Human Gene Id :	26959
Human Swiss Prot	O60381
No:	
Mouse Gene Id :	73389
Mouse Swiss Prot No :	Q8R316
Rat Gene Id :	27080
Rat Swiss Prot No :	Q62661
Immunogen :	The antiserum was produced against synthesized peptide derived from human HBP1. AA range:431-480
Specificity :	HBP1 Polyclonal Antibody detects endogenous levels of HBP1 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:10000 IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity- chromatography using epitope-specific immunogen.



Best Tools for immunology Research		
Concentration :	1 mg/ml	
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)	
Observed Band :	57kD	
Background :	function:Transcriptional repressor that binds to the promoter region of target genes. Plays a role in the regulation of the cell cycle and of the Wnt pathway. Binds preferentially to the sequence 5'-TTCATTCATTCA-3'. Binding to the H1F0 promoter is enhanced by interaction with RB1. Disrupts the interaction between DNA and TCF4.,similarity:Contains 1 AXH domain.,similarity:Contains 1 HMG box DNA-binding domain.,subunit:Binds the second PAH repeat of SIN3A (By similarity). Binds TCF4 and RB1.,	
Function :	function:Transcriptional repressor that binds to the promoter region of target genes. Plays a role in the regulation of the cell cycle and of the Wnt pathway. Binds preferentially to the sequence 5'-TTCATTCATTCA-3'. Binding to the H1F0 promoter is enhanced by interaction with RB1. Disrupts the interaction between DNA and TCF4.,similarity:Contains 1 AXH domain.,similarity:Contains 1 HMG box DNA-binding domain.,subunit:Binds the second PAH repeat of SIN3A (By similarity). Binds TCF4 and RB1.,	
Subcellular	Nucleus .	
Location : Expression :	Brain,Hepatoma,Uterus,	
Tag :	hot	
Sort :	7257	
No4 :		

## Products Images







Immunohistochemical analysis of paraffin-embedded human Colon cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).