


MKP-3 Polyclonal Antibody


Catalog No :	YT2774
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC;IF;ELISA
Target :	MKP-3
Fields :	>>MAPK signaling pathway;>>Transcriptional misregulation in cancer;>>Acute myeloid leukemia
Gene Name :	DUSP6
Protein Name :	Dual specificity protein phosphatase 6
Human Gene Id :	1848
Human Swiss Prot No :	Q16828
Mouse Gene Id :	67603
Mouse Swiss Prot No :	Q9DBB1
Rat Gene Id :	116663
Rat Swiss Prot No :	Q64346
Immunogen :	The antiserum was produced against synthesized peptide derived from human DUSP6. AA range:61-110
Specificity :	MKP-3 Polyclonal Antibody detects endogenous levels of MKP-3 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:20000. Not yet tested in other applications.

Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	42kD
Cell Pathway :	MAPK_ERK_Growth;MAPK_G_Protein;
Background :	The protein encoded by this gene is a member of the dual specificity protein phosphatase subfamily. These phosphatases inactivate their target kinases by dephosphorylating both the phosphoserine/threonine and phosphotyrosine residues. They negatively regulate members of the mitogen-activated protein (MAP) kinase superfamily (MAPK/ERK, SAPK/JNK, p38), which are associated with cellular proliferation and differentiation. Different members of the family of dual specificity phosphatases show distinct substrate specificities for various MAP kinases, different tissue distribution and subcellular localization, and different modes of inducibility of their expression by extracellular stimuli. This gene product inactivates ERK2, is expressed in a variety of tissues with the highest levels in heart and pancreas, and unlike most other members of this family, is localized in the cytoplasm. Mutations in t
Function :	catalytic activity:A phosphoprotein + H(2)O = a protein + phosphate.,catalytic activity:Protein tyrosine phosphate + H(2)O = protein tyrosine + phosphate.,function:Inactivates MAP kinases. Has a specificity for the ERK family.,similarity:Belongs to the protein-tyrosine phosphatase family. Non-receptor class dual specificity subfamily.,similarity:Contains 1 rhodanese domain.,similarity:Contains 1 tyrosine-protein phosphatase domain.,
Subcellular Location :	Cytoplasm .
Expression :	Expressed in keratinocytes (at protein level).
Sort :	9664
No4 :	1


Products Images



Western Blot analysis of HEPG2-UV cells using MKP-3
Polyclonal Antibody diluted at 1:500



Immunofluorescence analysis of HeLa cells, using DUSP6
Antibody. The picture on the right is blocked with the synthesized
peptide.



Immunohistochemistry analysis of paraffin-embedded human
brain tissue, using DUSP6 Antibody. The picture on the right is
blocked with the synthesized peptide.