

PFK-2 car Polyclonal Antibody

Catalog No: YT3681

Reactivity: Human; Mouse; Rat

Applications: WB;IHC;IF;ELISA

Target: PFK-2 car

Fields: >>Fructose and mannose metabolism;>>Metabolic pathways;>>AMPK

signaling pathway;>>Thyroid hormone signaling pathway

Gene Name: PFKFB2

Protein Name: 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase 2

O60825

P70265

Human Gene Id: 5208

Human Swiss Prot

No:

Mouse Gene Id: 18640

Mouse Swiss Prot

No:

Rat Gene Id: 24640

Rat Swiss Prot No: Q9JJH5

Immunogen : The antiserum was produced against synthesized peptide derived from human

PFKFB2. AA range:451-500

Specificity: PFK-2 car Polyclonal Antibody detects endogenous levels of PFK-2 car protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution : WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:5000.. IF 1:50-200

1/3



Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 58kD

Cell Pathway: Fructose and mannose metabolism;

Background: The protein encoded by this gene is involved in both the synthesis and

degradation of fructose-2,6-bisphosphate, a regulatory molecule that controls glycolysis in eukaryotes. The encoded protein has a 6-phosphofructo-2-kinase activity that catalyzes the synthesis of fructose-2,6-bisphosphate, and a fructose-2,6-biphosphatase activity that catalyzes the degradation of

fructose-2,6-bisphosphate. This protein regulates fructose-2,6-bisphosphate levels in the heart, while a related enzyme encoded by a different gene regulates fructose-2,6-bisphosphate levels in the liver and muscle. This enzyme functions as a homodimer. Two transcript variants encoding two different isoforms have

been found for this gene. [provided by RefSeq, Jul 2008],

Function: catalytic activity:ATP + D-fructose 6-phosphate = ADP + beta-D-fructose

2,6-bisphosphate.,catalytic activity:Beta-D-fructose 2,6-bisphosphate + H(2)O = D-fructose 6-phosphate + phosphate.,enzyme regulation:Phosphorylation results in the activation of the kinase activity.,function:Synthesis and degradation of fructose 2,6-bisphosphate.,similarity:In the C-terminal section; belongs to the phosphoglycerate mutase family.,subunit:Homodimer.,tissue specificity:Heart.,

Subcellular Location:

cytosol,

Heart.

Expression:

Tag:

orthogonal,hot

Sort:

11853


Host:

Rabbit


Modifications:

Unmodified


Products Images



Western Blot analysis of various cells using PFK-2 car Polyclonal Antibody



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue, using PFKFB2 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from 293 cells, treated with Heat shock, using PFKFB2 Antibody. The lane on the right is blocked with the synthesized peptide.