

Rab 18 Polyclonal Antibody

Catalog No: YT3920

Reactivity: Human; Mouse; Rat

Applications: WB;ELISA;IHC

Target: Rab 18

Gene Name: RAB18

Protein Name: Ras-related protein Rab-18

Human Gene Id: 22931

Human Swiss Prot

No:

Q9NP72

P35293

--

Mouse Gene ld: 19330

Mouse Swiss Prot

No:

Rat Gene Id: 307039

Rat Swiss Prot No: Q5EB77

Immunogen: The antiserum was produced against synthesized peptide derived from human

RAB18. AA range:103-152

Specificity: Rab 18 Polyclonal Antibody detects endogenous levels of Rab 18 protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1:500-2000;IHC 1:50-300; ELISA 2000-20000

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.



Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 28kD

Background: The protein encoded by this gene is a member of a family of Ras-related small

GTPases that regulate membrane trafficking in organelles and transport vesicles. Knockdown studies is zebrafish suggest that this protein may have a role in eye and brain development. Mutations in this gene are associated with Warburg micro syndrome type 3. Alternatively spliced transcript variants have been found for this

gene. [provided by RefSeq, Jan 2012],

Function: function:Plays a role in apical endocytosis/recycling. May be implicated in

transport between the plasma membrane and early

endosomes., similarity: Belongs to the small GTPase superfamily. Rab

family., tissue specificity: Ubiquitous.,

Subcellular Location:

Apical cell membrane . Lipid droplet .


Expression: Ubiquitous.

Sort : 13211


Host: Rabbit

Modifications: Unmodified


Products Images




Western Blot analysis of MCF-7 cells using Rab 18 Polyclonal Antibody



Western blot analysis of lysates from MCF-7 and HUVEC cells, using RAB18 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HepG2 cells using RAB18 antibody.



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).