

RASL10B Polyclonal Antibody

| Catalog No : | YT4010 | | |
|-------------------------|---|--|--|
| Reactivity : | Human;Mouse | | |
| Applications : | WB;IHC | | |
| Target : | RASL10B | | |
| Gene Name : | RASL10B | | |
| Protein Name : | Ras-like protein family member 10B | | |
| Human Gene Id : | 91608 | | |
| Human Swiss Prot | Q96S79 | | |
| No : Mouse Gene Id : | 276952 | | |
| Mouse Swiss Prot | Q5SSG5 | | |
| No : Immunogen : | Synthesized peptide derived from RASL10B . at AA range: 80-160 | | |
| Specificity : | RASL10B Polyclonal Antibody detects endogenous levels of RASL10B protein. | | |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. | | |
| Source : | Polyclonal, Rabbit,IgG | | |
| Dilution : | WB 1:500-2000;IHC 1:50-300 | | |
| Purification : | The antibody was affinity-purified from rabbit antiserum by affinity- chromatography using epitope-specific immunogen. | | |
| Concentration : | 1 mg/ml | | |
| Storage Stability : | -15°C to -25°C/1 year(Do not lower than -25°C) | | |




Observed Band : 22kD

| Background : | similarity:Belongs to the small GTPase superfamily. Ras family., | | |
|-----------------|--|--|--|
| | | | |
| Function : | similarity:Belongs to the small GTPase superfamily. Ras family., | | |
| Subcellular | Cell membrane ; Lipid-anchor ; Cytoplasmic side . | | |
| Expression : | Expressed at high levels in skeletal muscle and, at much lower levels, in heart, brain and pancreas. | | |
| Sort : | 13827 | | |
| Soft. | 13027 | | |
| No4 : | 1 | | |
| Host : | Rabbit | | |
| Modifications : | Unmodified | | |

| Products Images | | | | |
|-----------------|--|---|--|--|
| 31 | 3 250 150 100 75 50 37 | Western blot analysis of RASL10B Antibody. The lane on the right is blocked with the RASL10B peptide. | | |
| RASL10B | 25 20 | | | |
| | 15 (kd) | | | |





Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).