


Synaptotagmin XIII Polyclonal Antibody

Catalog No :	YT4488
Reactivity :	Human;Mouse;Rat
Applications :	WB;ELISA
Target :	Synaptotagmin XIII
Gene Name :	SYT13
Protein Name :	Synaptotagmin-13
Human Gene Id :	57586
Human Swiss Prot No :	Q7L8C5
Mouse Gene Id :	80976
Mouse Swiss Prot No :	Q9EQT6
Rat Gene Id :	80977
Rat Swiss Prot No :	Q925B5
Immunogen :	The antiserum was produced against synthesized peptide derived from human SYT13. AA range:321-370
Specificity :	Synaptotagmin XIII Polyclonal Antibody detects endogenous levels of Synaptotagmin XIII protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	48kD
Background :	<p>synaptotagmin 13(SYT13) Homo sapiens This gene encodes a member of the large synaptotagmin protein family. Family members have an extracellular N-terminal transmembrane domain and a cytoplasmic C terminus with two tandem C2 domains (C2A and C2B). Synaptotagmin family members can form homo- and heteromeric complexes with each other. They also have different biochemical properties and developmental profiles, and patterns of tissue distribution. Synaptotagmins function as membrane traffickers in multicellular organisms. Two alternatively spliced transcript variants that encode different protein isoforms have been described for this gene. [provided by RefSeq, Oct 2011],</p>
Function :	<p>domain:The first C2 domain/C2A does not mediate Ca(2+)-dependent phospholipid binding.,domain:The second C2 domain/C2B domain binds phospholipids regardless of whether calcium is present.,function:May be involved in transport vesicle docking to the plasma membrane.,similarity:Belongs to the synaptotagmin family.,similarity:Contains 2 C2 domains.,subunit:Interacts with NRXN1.,tissue specificity:Expressed in brain, pancreas and kidney.,</p>
Subcellular Location :	Membrane ; Single-pass membrane protein .
Expression :	Expressed in brain, pancreas and kidney.
Sort :	16815
No4 :	1

Products Images



Western Blot analysis of various cells using Synaptotagmin XIII Polyclonal Antibody



Western blot analysis of SYT13 Antibody. The lane on the right is blocked with the SYT13 peptide.