

TALK-2 Polyclonal Antibody

Catalog No: YT4539

Reactivity: Human; Rat; Mouse;

Applications: WB;ELISA

Target: TALK-2

Gene Name: KCNK17

Protein Name: Potassium channel subfamily K member 17

Q96T54

Human Gene Id: 89822

Human Swiss Prot

No:

Immunogen: The antiserum was produced against synthesized peptide derived from human

KCNK17. AA range:271-320

Specificity: TALK-2 Polyclonal Antibody detects endogenous levels of TALK-2 protein.

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1:500 - 1:2000. ELISA: 1:20000. Not yet tested in other applications.

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 37-42kD

Background: potassium two pore domain channel subfamily K member 17(KCNK17) Homo

sapiens The protein encoded by this gene belongs to the family of potassium



channel proteins containing two pore-forming P domains. This channel is an open rectifier which primarily passes outward current under physiological K+ concentrations. This gene is activated at alkaline pH. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2008],

Function:

function:Outward rectifying potassium channel. Produces rapidly activating and non-inactivating outward rectifier K(+) currents.,miscellaneous:Inhibited by Ba(2+), quinidine, chloroform and halothane. Activated at alkaline pH. Activated by quinine and isoflurane.,similarity:Belongs to the two pore domain potassium channel (TC 1.A.1.8) family.,subunit:Homodimer .,

Subcellular Location:


Membrane; Multi-pass membrane protein.

Expression: Adrenal gland, Lung, Pancreas, Spleen,


Sort : 16907

No4:

Products Images



Immunofluorescence analysis of Caco2 cell. 1,primary Antibody was diluted at 1:100(4°C overnight). 2, Goat Anti Rabbit IgG (H&L) - AFluor 594 Secondary antibody(catalog No: RS3611) was diluted at 1:500(room temperature, 50min).



Western blot analysis of lysates from Jurkat cells, using KCNK17 Antibody. The lane on the right is blocked with the synthesized peptide.