

## v-SNARE Vti1p Polyclonal Antibody


Catalog No :	YT4895
Reactivity :	Human;Mouse;Rat
Applications :	WB;ELISA
Target :	v-SNARE Vti1p
Fields :	>>SNARE interactions in vesicular transport
Gene Name :	VTI1B
Protein Name :	Vesicle transport through interaction with t-SNAREs homolog 1B
Human Gene Id :	10490
Human Swiss Prot	Q9UEU0
No :	
Mouse Swiss Prot No :	O88384
Rat Swiss Prot No :	P58200
Immunogen :	Synthesized peptide derived from the Internal region of human v-SNARE Vti1p.
Specificity :	v-SNARE Vti1p Polyclonal Antibody detects endogenous levels of v-SNARE Vti1p protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity- chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml



Best rools for initiatiology Research	
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	26kD
Cell Pathway :	SNARE interactions in vesicular transport;
Background :	function:V-SNARE that mediates vesicle transport pathways through interactions with t-SNAREs on the target membrane. These interactions are proposed to mediate aspects of the specificity of vesicle trafficking and to promote fusion of the lipid bilayers. May be concerned with increased secretion of cytokines associated with cellular senescence.,similarity:Belongs to the VTI1 family.,subunit:Found in a complex with VAMP8 and STX7 in the liver. Forms a SNARE complex with STX7, STX8 and VAMP8 which functions in the homotypic fusion of late endosomes. Part of the SNARE core complex containing STX7, STX8 and VAMP8.,tissue specificity:Expressed in all tissues examined.,
Function :	function:V-SNARE that mediates vesicle transport pathways through interactions with t-SNAREs on the target membrane. These interactions are proposed to mediate aspects of the specificity of vesicle trafficking and to promote fusion of the lipid bilayers. May be concerned with increased secretion of cytokines associated with cellular senescence.,similarity:Belongs to the VTI1 family.,subunit:Found in a complex with VAMP8 and STX7 in the liver. Forms a SNARE complex with STX7, STX8 and VAMP8 which functions in the homotypic fusion of late endosomes. Part of the SNARE core complex containing STX7, STX8 and VAMP8.,tissue specificity:Expressed in all tissues examined.,
Subcellular Location :	Early endosome membrane ; Single-pass type IV membrane protein . Late endosome membrane ; Single-pass type IV membrane protein . Lysosome membrane . Cytoplasmic granule . Recycling endosome membrane ; Single-pass type IV membrane protein .
Expression :	Expressed in all tissues examined.
Sort :	24216
No4 :	1
Host :	Rabbit
Modifications :	Unmodified

## Products Images





Western Blot analysis of RAT-HEART cells using v-SNARE Vti1p Polyclonal Antibody diluted at 1:1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000