

PDP2 Polyclonal Antibody

Catalog No: YT5074

Reactivity: Human; Rat; Mouse;

Applications: WB;ELISA;IHC

Target: PDP2

Gene Name: PDP2

Protein Name: [Pyruvate dehydrogenase [acetyl-transferring]]-phosphatase 2 mitochondrial

Human Gene Id: 57546

Human Swiss Prot

No:

Immunogen: Synthesized peptide derived from PDP2 . at AA range: 70-150

Specificity: PDP2 Polyclonal Antibody detects endogenous levels of PDP2 protein.

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Q9P2J9

Dilution: WB 1:500-2000;IHC 1:50-300; ELISA 2000-20000

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 59kD

Background: This gene is a mitochondrial protein that functions as a phosphatase and is

involved in the enzymatic resetting of the pyruvate dehydrogenase complex. Alternative splicing results in multiple transcript variants encoding the same



protein. [provided by RefSeq, Aug 2016],

Function: catalytic activity:[Pyruvate dehydrogenase (acetyl-transferring)] phosphate +

H(2)O = [pyruvate dehydrogenase (acetyl-transferring)] +

phosphate.,cofactor:Binds 2 magnesium ions per subunit.,function:Catalyzes the dephosphorylation and concomitant reactivation of the alpha subunit of the E1 component of the pyruvate dehydrogenase complex.,similarity:Belongs to the PP2C family.,subunit:Heterodimer of a catalytic subunit and a FAD protein of

unknown function.,

Subcellular Location :

Mitochondrion matrix.


Expression: Brain,

Sort : 11791


Host: Rabbit

Modifications: Unmodified

Products Images



Western Blot analysis of extracts from K562 cells, using PDP2 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).