

CANB2 Polyclonal Antibody

Catalog No: YN0515

Reactivity: Human;Rat

Applications: WB;ELISA

Target: CANB2

Fields: >>MAPK signaling pathway;>>Calcium signaling pathway;>>cGMP-PKG

signaling pathway;>>Oocyte meiosis;>>Cellular senescence;>>Wnt signaling

pathway;>>Axon guidance;>>VEGF signaling pathway;>>Osteoclast

differentiation;>>C-type lectin receptor signaling pathway;>>Natural killer cell

mediated cytotoxicity;>>Th1 and Th2 cell differentiation;>>Th17 cell

differentiation;>>T cell receptor signaling pathway;>>B cell receptor signaling pathway;>>Long-term potentiation;>>Glutamatergic synapse;>>Oxytocin

signaling pathway;>>Glucagon signaling pathway;>>Renin

secretion;>>Alzheimer disease;>>Amyotrophic lateral sclerosis;>>Prion

disease;>>Pathways of neurodegeneration - multiple diseases;>>Amphetamine addiction:>>Tuberculosis:>>Human cytomegalovirus infection:>>Human T-cell

leukemia virus 1 infection;>>Kaposi sarcoma-associated herpesvirus

infection;>>Human immunodeficiency virus 1 infection;>>PD-L1 expression and

PD-1 checkpoint pathway in cancer;>>Lipid and atherosclerosis

Gene Name: PPP3R2 CBLP PPP3RL

Protein Name: Calcineurin subunit B type 2 (Calcineurin B-like protein) (CBLP) (Calcineurin BII)

(CNBII) (PPP3R1-like) (Protein phosphatase 2B regulatory subunit 2) (Protein

phosphatase 3 regulatory subunit B beta

Human Gene Id: 5535

Human Swiss Prot

Q96LZ3

No:

Mouse Swiss Prot

Q63811

No:

Rat Swiss Prot No: P28470

Immunogen: Synthesized peptide derived from part region of human protein

Specificity: CANB2 Polyclonal Antibody detects endogenous levels of protein.



Host:


Rabbit

Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Formulation: Source: Polyclonal, Rabbit, IgG **Dilution:** WB 1:500-2000 ELISA 1:5000-20000 **Purification:** The antibody was affinity-purified from rabbit antiserum by affinitychromatography using epitope-specific immunogen. 1 mg/ml Concentration: -15°C to -25°C/1 year(Do not lower than -25°C) Storage Stability: **Observed Band:** 18kD MAPK_ERK_Growth;MAPK_G_Protein;Calcium;Oocyte meiosis;Apoptosis_Inhi **Cell Pathway:** bition: Apoptosis Mitochondrial: Apoptosis Overview: WNT: WNT-T CELLAxon guidance; VEGF; Natural killer cell mediated cytotoxicity; T Cell function: Regulatory subunit of calcineurin, a calcium-dependent, calmodulin **Background:** stimulated protein phosphatase. Confers calcium sensitivity..miscellaneous:This protein has four functional calcium-binding sites., similarity: Belongs to the calcineurin regulatory subunit family., similarity: Contains 4 EF-hand domains., subunit: Composed of a catalytic subunit (A) and a regulatory subunit (B).,tissue specificity:Testis specific., **Function:** function: Regulatory subunit of calcineurin, a calcium-dependent, calmodulin stimulated protein phosphatase. Confers calcium sensitivity., miscellaneous: This protein has four functional calcium-binding sites., similarity: Belongs to the calcineurin regulatory subunit family., similarity: Contains 4 EF-hand domains., subunit: Composed of a catalytic subunit (A) and a regulatory subunit (B).,tissue specificity:Testis specific., Subcellular Mitochondrion. Localizes in the mitochondria in a SPATA33-dependent manner. Location: **Expression:** Testis-specific. Sort: 18202 No4:



Modifications: Unmodified

Products Images



Western blot analysis of lysates from KB cells, primary antibody was diluted at 1:1000, 4° over night