

## ROR1 rabbit pAb

YT7999 **Catalog No:** 

Human; Mouse; Rat Reactivity:

**Applications:** WB

Target: ROR1

Fields: >>Wnt signaling pathway

Q01973

Q9Z139

**Gene Name: ROR1 NTRKR1** 

**Protein Name:** ROR1

**Human Gene Id:** 4919

**Human Swiss Prot** 

No:

Mouse Gene Id: 26563

**Mouse Swiss Prot** 

No:

Synthesized peptide derived from human ROR1 Immunogen:

This antibody detects endogenous levels of ROR1 at Human, Mouse, Rat **Specificity:** 

**Formulation:** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

**Dilution:** WB 1:500-2000

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Concentration:** 1 mg/ml

1/2



Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 105kD

Location:

**Background :** This gene encodes a receptor tyrosine kinase-like orphan receptor that

modulates neurite growth in the central nervous system. The encoded protein is a glycosylated type I membrane protein that belongs to the ROR subfamily of cell surface receptors. It is a pseudokinase that lacks catalytic activity and may interact with the non-canonical Wnt signalling pathway. This gene is highly expressed during early embryonic development but expressed at very low levels in adult tissues. Increased expression of this gene is associated with B-cell chronic lymphocytic leukaemia. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jun 2012],

**Function :** catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine

phosphate., developmental stage: Expressed at high levels during early embryonic development. The expression levels drop strongly around day 16 and there are only very low levels in adult tissues., function: Tyrosine-protein kinase receptor whose role is not yet clear., similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family. ROR subfamily., similarity: Contains 1 FZ (frizzled)

domain., similarity: Contains 1 Ig-like C2-type (immunoglobulin-like)

domain., similarity: Contains 1 kringle domain., similarity: Contains 1 protein kinase domain., tissue specificity: Expressed strongly in human heart, lung, and kidney, but weakly in the CNS. The short isoform is strongly expressed in fetal and adult CNS and in a variety of human cancers, including those originating from CNS or

PNS neuroectoderm.,

Subcellular Membrane ; Single-pass type I membrane protein. Cell projection, axon .

**Expression:** Expressed strongly in human heart, lung and kidney, but weakly in the CNS.

Isoform Short is strongly expressed in fetal and adult CNS and in a variety of

human cancers, including those originating from CNS or PNS neuroectoderm.

**Sort :** 25072

**No4:** 1

## **Products Images**

2/2