

## Retinoic Acid Receptor $\alpha/\beta$ rabbit pAb

<b>Catalog No :</b>	YT8037
<b>Reactivity :</b>	Human;Mouse;Rat
<b>Applications :</b>	WB
<b>Target :</b>	Retinoic Acid Receptor $\alpha/\beta$
<b>Fields :</b>	>>Th17 cell differentiation;>>Estrogen signaling pathway;>>Pathways in cancer;>>Transcriptional misregulation in cancer;>>Acute myeloid leukemia
<b>Gene Name :</b>	RARA NR1B1
<b>Protein Name :</b>	Retinoic Acid Receptor $\alpha/\beta$
<b>Human Gene Id :</b>	5914
<b>Human Swiss Prot No :</b>	P10276
<b>Mouse Gene Id :</b>	19401
<b>Mouse Swiss Prot No :</b>	P11416
<b>Immunogen :</b>	Synthesized peptide derived from human Retinoic Acid Receptor $\alpha/\beta$
<b>Specificity :</b>	This antibody detects endogenous levels of Retinoic Acid Receptor $\alpha/\beta$ at Human, Mouse,Rat
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500-2000
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml

**Storage Stability :** -15°C to -25°C/1 year (Do not lower than -25°C)

**Observed Band :** 50kD

**Background :** This gene represents a nuclear retinoic acid receptor. The encoded protein, retinoic acid receptor alpha, regulates transcription in a ligand-dependent manner. This gene has been implicated in regulation of development, differentiation, apoptosis, granulopoiesis, and transcription of clock genes. Translocations between this locus and several other loci have been associated with acute promyelocytic leukemia. Alternatively spliced transcript variants have been found for this locus. [provided by RefSeq, Sep 2010],

**Function :** disease: Chromosomal aberrations involving RARA may be a cause of acute promyelocytic leukemia (APL) [MIM:612376]. Translocation t(11;17)(q32;q21) with ZBTB16/PLZF; translocation t(15;17)(q21;q21) with PML; translocation t(5;17)(q32;q11) with NPM., domain: Composed of three domains: a modulating N-terminal domain, a DNA-binding domain and a C-terminal steroid-binding domain., function: This is a receptor for retinoic acid. This metabolite has profound effects on vertebrate development. Retinoic acid is a morphogen and is a powerful teratogen. This receptor controls cell function by directly regulating gene expression., online information: Retinoic acid receptor entry, PTM: Phosphorylated. Phosphorylation does not change during cell cycle. Phosphorylation on Ser-77 is crucial for transcriptional activity., similarity: Belongs to the nuclear hormone receptor family., similarity: Belongs to the nuclear

**Subcellular Location :** Nucleus . Cytoplasm . Nuclear localization depends on ligand binding, phosphorylation and sumoylation (PubMed:19850744). Translocation to the nucleus in the absence of ligand is dependent on activation of PKC and the downstream MAPK phosphorylation (By similarity). Increased nuclear localization upon pulsatile shear stress (PubMed:28167758). .

**Expression :** Expressed in monocytes.

**Sort :** 25108

**No4 :** 1

**Products Images**