

## MAPKAPK-2 (phospho Thr222) Polyclonal Antibody

Catalog No :	YP0692
Reactivity :	Human;Mouse;Rat;Monkey
Applications :	WB;IHC;IF;ELISA
Target :	МАРКАРК2
Fields :	>>MAPK signaling pathway;>>Cellular senescence;>>VEGF signaling pathway;>>C-type lectin receptor signaling pathway;>>Neurotrophin signaling pathway;>>Kaposi sarcoma-associated herpesvirus infection;>>Viral carcinogenesis
Gene Name :	MAPKAPK2
Protein Name :	MAP kinase-activated protein kinase 2
Human Gene Id :	9261
Human Swiss Prot	P49137
No : Mouse Gene Id :	17164
Mouse Swiss Prot	P49138
No : Immunogen :	The antiserum was produced against synthesized peptide derived from human MAPKAPK-2 around the phosphorylation site of Thr222. AA range:188-237
Specificity :	Phospho-MAPKAPK-2 (T222) Polyclonal Antibody detects endogenous levels of MAPKAPK-2 protein only when phosphorylated at T222.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:20000 IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-



chromatography using epitope-specific immunogen.

Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	45kD
Cell Pathway :	MAPK_ERK_Growth;MAPK_G_Protein;VEGF;Neurotrophin;
Background :	This gene encodes a member of the Ser/Thr protein kinase family. This kinase is regulated through direct phosphorylation by p38 MAP kinase. In conjunction with p38 MAP kinase, this kinase is known to be involved in many cellular processes including stress and inflammatory responses, nuclear export, gene expression regulation and cell proliferation. Heat shock protein HSP27 was shown to be one of the substrates of this kinase in vivo. Two transcript variants encoding two different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],
Function :	catalytic activity:ATP + a protein = ADP + a phosphoprotein.,enzyme regulation:Seems to be activated by two distinct pathways: the first involves the stimulation of p42/p44 MAPK by growth factors, the second, triggered by stress and heat shock, depends on the activation of MPK2 and upstream MAPKK/MAPKKK.,function:Its physiological substrate seems to be the small heat shock protein (HSP27/HSP25). In vitro can phosphorylate glycogen synthase at 'Ser-7' and tyrosine hydroxylase (on 'Ser-19' and 'Ser-40'). This kinase phosphorylates Ser in the peptide sequence, Hyd-X-R-X(2)-S, where Hyd is a large hydrophobic residue (By similarity). Mediates both ERK and p38 MAPK/MAPK14 dependent neutrophil responses. Participates in TNF alpha- stimulated exocytosis of secretory vesicles in neutrophils. Plays a role in phagocytosis-induced respiratory burst activity.,PTM:Phosphorylated and activated by MAP k
Subcellular Location :	Cytoplasm . Nucleus . Phosphorylation and subsequent activation releases the autoinhibitory helix, resulting in the export from the nucleus into the cytoplasm.
Expression :	Expressed in all tissues examined.
Tag :	orthogonal,hot
Sort :	9376
No2 :	3316T
No4 :	1



0.700

0.600

0.500

0.400 0.300 0.200 0.100

0.000



0.061

non-phosphopeptide

## **Products Images**

Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using MAPKAPK-2 (Phospho-Thr222) Antibody



OD 450nm Reading

phosphopeptide

Immunohistochemistry analysis of paraffin-embedded human heart, using MAPKAPK-2 (Phospho-Thr222) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from COS7 cells treated with UV 15', using MAPKAPK-2 (Phospho-Thr222) Antibody. The lane on the right is blocked with the phospho peptide.