

FRS2 (phospho Tyr436) Polyclonal Antibody

Catalog No: YP0805

Reactivity: Human; Mouse; Monkey

Applications: WB;IHC;IF;ELISA

Target: FRS2

Fields: >>Thermogenesis;>>Neurotrophin signaling pathway;>>Proteoglycans in

cancer

Gene Name: FRS2

Protein Name: Fibroblast growth factor receptor substrate 2

Q8WU20

Q8C180

Human Gene Id: 10818

Human Swiss Prot

No:

Mouse Gene ld: 327826

Mouse Swiss Prot

No:

Immunogen: The antiserum was produced against synthesized peptide derived from human

FRS2 around the phosphorylation site of Tyr436. AA range:402-451

Specificity: Phospho-FRS2 (Y436) Polyclonal Antibody detects endogenous levels of FRS2

protein only when phosphorylated at Y436.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:10000.. IF 1:50-200

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.



Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 65kD

Cell Pathway: Neurotrophin;

Background: function: Adapter protein that links FGR and NGF receptors to downstream

signaling pathways. Involved in the activation of MAP kinases. Modulates

signaling via SHC1 by competing for a common binding site on

NTRK1.,PTM:Phosphorylated on tyrosine residues upon stimulation by NGF.,PTM:Ubiquitinated when tyrosine phosphorylated and in a complex with GRB2. The unphosphorylated form is not subject to ubiquitination.,sequence

caution: Translated as stop., similarity: Contains 1 IRS-type PTB

domain.,subcellular location:Cytoplasmic, membrane-bound.,subunit:Part of a complex containing FRS2, GRB2 and SOS1. Part of a complex containing GRB2 and CBL. Binds RET (By similarity). Binds FGFR1, SUC1, NTRK1, NTRK2, NTRK3 and SRC. The tyrosine-phosphorylated protein binds the SH2 domains of GRB2 and PTPN11.,tissue specificity:Highly expressed in heart, brain, spleen,

lung, liver, skeletal muscle, kidney and testis.,

Function: function: Adapter protein that links FGR and NGF receptors to downstream

signaling pathways. Involved in the activation of MAP kinases. Modulates

signaling via SHC1 by competing for a common binding site on

NTRK1.,PTM:Phosphorylated on tyrosine residues upon stimulation by NGF.,PTM:Ubiquitinated when tyrosine phosphorylated and in a complex with GRB2. The unphosphorylated form is not subject to ubiquitination.,sequence

caution:Translated as stop., similarity:Contains 1 IRS-type PTB

domain.,subcellular location:Cytoplasmic, membrane-bound.,subunit:Part of a complex containing FRS2, GRB2 and SOS1. Part of a complex containing GRB2 and CBL. Binds RET (By similarity). Binds FGFR1, SUC1, NTRK1, NTRK2, NTRK3 and SRC. The tyrosine-phosphorylated protein binds the SH2 domains of GRB2 and PTPN11.,tissue specificity:Highly expressed in heart, brain, spleen,

lung, liver, skeletal muscle, kidney and t

Subcellular Endomembrane system. Cytoplasmic, membrane-bound.

Location:

Expression: Highly expressed in heart, brain, spleen, lung, liver, skeletal muscle, kidney and

testis.

Tag: orthogonal

Sort: 6306

No2: 3861S


2/4

1


No4:

Products Images


Western Blot analysis of various cells using Phospho-FRS2 (Y436) Polyclonal Antibody




- + phospho-peptide
- + non-phospho-peptide
- + + + 3T3 NGF(customer's sample)



Western Blot analysis of COS cells using Phospho-FRS2 (Y436) Polyclonal Antibody



Immunohistochemistry analysis of paraffin-embedded human brain, using FRS2 (Phospho-Tyr436) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from COS7 cells, using FRS2 (Phospho-Tyr436) Antibody. The lane on the right is blocked with the phospho peptide.