

TAL1/2 (Acetyl Lys221/Acetyl Lys222/Acetyl Lys36/Acetyl Lys37) Polyclonal Antibody

Catalog No :	YK0071
Reactivity :	Human;Mouse;Rat
Applications :	WB;ELISA
Target :	TAL1/2
Gene Name :	TAL1/TAL2
Protein Name :	T-cell acute lymphocytic leukemia protein 1 homolog/T-cell acute lymphocytic leukemia protein 2
Human Gene Id :	6886
Human Swiss Prot No :	P17542
Mouse Gene Id :	21349
Mouse Swiss Prot No :	P22091
Immunogen :	Synthesized acetyl-peptide derived from human TAL1/2 around the acetylation site of K221.
Specificity :	Acetyl-TAL1/2 (K221/K222/K36/K37) Polyclonal Antibody detects endogenous levels of TAL1/2 around the acetylation site of K221 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml

Storage Stability : -15°C to -25°C/1 year (Do not lower than -25°C)

Observed Band : 45kD

Background : alternative products: The splicing pattern is cell-lineage dependent, disease: A chromosomal aberration involving TAL1 may be a cause of some T-cell acute lymphoblastic leukemias (T-ALL). Translocation t(1;14)(p32;q11) with T-cell receptor alpha chain (TCRA) genes., domain: The helix-loop-helix domain is necessary and sufficient for the interaction with DRG1., function: Implicated in the genesis of hemopoietic malignancies. It may play an important role in hemopoietic differentiation. Serves as a positive regulator of erythroid differentiation., PTM: Phosphorylated on serine residues. Phosphorylation of Ser-122 is strongly stimulated by hypoxia., PTM: Ubiquitinated; subsequent to hypoxia-dependent phosphorylation of Ser-122, ubiquitination targets the protein for rapid degradation via the ubiquitin system. This process may be characteristic for microvascular endothelial cells, since it could not be observed in large vessel endothelial cells., similarity: Contains 1 basic helix-loop-helix (bHLH) domain., subunit: Efficient DNA binding requires dimerization with another bHLH protein. Forms heterodimers with TCF3. Binds to the LIM domain containing protein LMO2 and to DRG1. Can assemble in a complex with LDB1 and LMO2. Component of a TAL-1 complex composed at least of CBFA2T3, LDB1, TAL1 and TCF3., tissue specificity: Leukemic stem cell.,

Function : alternative products: The splicing pattern is cell-lineage dependent, disease: A chromosomal aberration involving TAL1 may be a cause of some T-cell acute lymphoblastic leukemias (T-ALL). Translocation t(1;14)(p32;q11) with T-cell receptor alpha chain (TCRA) genes., domain: The helix-loop-helix domain is necessary and sufficient for the interaction with DRG1., function: Implicated in the genesis of hemopoietic malignancies. It may play an important role in hemopoietic differentiation. Serves as a positive regulator of erythroid differentiation., PTM: Phosphorylated on serine residues. Phosphorylation of Ser-122 is strongly stimulated by hypoxia., PTM: Ubiquitinated; subsequent to hypoxia-dependent phosphorylation of Ser-122, ubiquitination targets the protein for rapid degradation via the ubiquitin system. This process may be characteristic for microvascular endothelial cells, since it could not be

Subcellular Location : Nucleus .

Expression : Leukemic stem cell.


Tag : orthogonal, hot

Sort : 16903

Host : Rabbit

Modifications : Acetyl

Products Images



Western blot analysis of MOUSE-HEART MOUSE-BRAIN using Acetyl-TAL1/2 (K221/K222/K36/K37) antibody. Antibody was diluted at 1:500. Secondary antibody(catalog#:RS0002) was diluted at 1:20000