

## CD70 (PN0445) Nb-FC recombinant antibody

YA0474 Catalog No:

Reactivity: Human

**Applications: ELISA** 

**Target:** CD70

Gene Name: CD70 CD27L CD27LG TNFSF7

**Protein Name:** CD70 antigen (CD27 ligand) (CD27-L) (Tumor necrosis factor ligand

superfamily member 7) (CD antigen CD70)

**Human Gene Id:** 970

**Human Swiss Prot** 

No:

Immunogen: Purified recombinant Human CD70

P32970

This recombinant monoclonal antibody can detects endogenous levels of CD70 **Specificity:** 

protein.

Formulation: Phosphate-buffered solution

Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain, Source:

recombinantly produced from 293F cell

**Dilution:** ELISA 1:5000-100000

**Purification:** Recombinant Expression and Affinity purified

Please check the information on the tube **Concentration:** 

-15°C to -25°C/1 year(Avoid freeze / thaw cycles) **Storage Stability:** 

**Cell Pathway:** Cytokine-cytokine receptor interaction;

**Background:** The protein encoded by This gene is a cytokine that belongs to the tumor

1/2

necrosis factor (TNF) ligand family. This cytokine is a ligand for TNFRSF27/CD27. It is a surface antigen on activated, but not on resting, T and B lymphocytes. It induces proliferation of costimulated T cells, enhances the generation of cytolytic T cells, and contributes to T cell activation. This cytokine is also reported to play a role in regulating B-cell activation, cytotoxic function of natural killer cells, and immunoglobulin sythesis. [provided by RefSeq, Jul 2008]

**Function:** Cytokine that binds to CD27. Plays a role in T-cell activation. Induces the

> proliferation of costimulated T-cells and enhances the generation of cytolytic Tcells., similarity: Belongs to the tumor necrosis factor family., subunit: Homotrimer.,

Subcellular Location:

Membrane; Single-pass type II membrane protein.

**Expression:** B-cell, Kidney

Tag: recombinant

Sort: 3652

No1: 14-0701-85

No4:

## **Products Images**

