

## CD73 (PN0521) Nb-FC recombinant antibody

Catalog No: YA0497

Reactivity: Human

**Applications:** ELISA

Target: CD73

Gene Name: NT5E NT5 NTE

**Protein Name:** 5'-nucleotidase (5'-NT) (EC 3.1.3.5) (Ecto-5'-nucleotidase) (CD antigen CD73)

Human Gene Id: 4907

**Human Swiss Prot** 

No:

Immunogen: Purified recombinant Human CD73

P21589

**Specificity:** This recombinant monoclonal antibody can detects endogenous levels of CD73

protein.

**Formulation:** Phosphate-buffered solution

Source: Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain,

recombinantly produced from 293F cell

**Dilution:** ELISA 1:5000-100000

Purification: Recombinant Expression and Affinity purified

**Concentration:** Please check the information on the tube

Storage Stability: -15°C to -25°C/1 year(Avoid freeze / thaw cycles)

**Cell Pathway:** Purine metabolism;Pyrimidine metabolism;Nicotinate and nicotinamide

metabolism;

**Background:** The protein encoded by This gene is a plasma membrane protein that catalyzes

1/2



the conversion of extracellular nucleotides to membrane-permeable nucleosides. The encoded protein is used as a determinant of lymphocyte differentiation. Defects inThis gene can lead to the calcification of joints and arteries. Two transcript variants encoding different isoforms have been found forThis gene.[provided by RefSeq, Mar 2011]

**Function:** catalytic activity: A 5'-ribonucleotide + H(2)O = a ribonucleoside +

phosphate.,cofactor:Zinc.,disease:There is a decrease in the activity of NT5 in B-cell chronic lymphocytic leukemia.,Hydrolyzes extracellular nucleotides into membrane permeable nucleosides.,similarity:Belongs to the 5'-nucleotidase

family., subunit: Homodimer; disulfide-linked.,

Subcellular Location:

Cell membrane ; Lipid-anchor, GPI-anchor .

**Expression :** Leukocyte, Liver, Placenta, Skin

Tag: recombinant

**Sort :** 3661

No4:

## **Products Images**

2/2