


CD80 Monoclonal Antibody


Catalog No :	YM0136
Reactivity :	Human
Applications :	IHC;IF;ELISA
Target :	CD80
Fields :	>>Cell adhesion molecules;>>Toll-like receptor signaling pathway;>>Intestinal immune network for IgA production;>>Type I diabetes mellitus;>>Autoimmune thyroid disease;>>Systemic lupus erythematosus;>>Rheumatoid arthritis;>>Allograft rejection;>>Graft-versus-host disease;>>Viral myocarditis
Gene Name :	CD80
Protein Name :	T-lymphocyte activation antigen CD80
Human Gene Id :	941
Human Swiss Prot No :	P33681
Mouse Swiss Prot No :	Q00609
Immunogen :	Purified recombinant fragment of CD80 expressed in E. Coli.
Specificity :	CD80 Monoclonal Antibody detects endogenous levels of CD80 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Monoclonal, Mouse
Dilution :	IHC 1:200 - 1:1000. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other applications.
Purification :	Affinity purification
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Cell Pathway :	Cell adhesion molecules (CAMs);Toll_Like;Intestinal immune network for IgA production;Type I diabetes mellitus;Autoimmune thyroid disease;Systemic lupus erythematosus;Allograft rejection;Graft-versus-
P References :	1. Transplant Proc. 2008 Oct;40(8):2729-33. 2. Nat Med. 2007 Dec;13(12):1440-9.
Background :	The protein encoded by this gene is a membrane receptor that is activated by the binding of CD28 or CTLA-4. The activated protein induces T-cell proliferation and cytokine production. This protein can act as a receptor for adenovirus subgroup B and may play a role in lupus neuropathy. [provided by RefSeq, Aug 2011],
Function :	function:Involved in the costimulatory signal essential for T-lymphocyte activation. T-cell proliferation and cytokine production is induced by the binding of CD28 or CTLA-4 to this receptor.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 1 Ig-like V-type (immunoglobulin-like) domain.,subunit:Interacts with adenovirus subgroup B fiber proteins and acts as a receptor for these viruses.,tissue specificity:Expressed on activated B-cells, macrophages and dendritic cells.,
Subcellular Location :	Membrane; Single-pass type I membrane protein.
Expression :	Expressed on activated B-cells, macrophages and dendritic cells.
Tag :	hot
Sort :	3681
No4 :	1

Products Images



Immunohistochemistry analysis of paraffin-embedded human Tonsil tissues with AEC staining using CD80 Monoclonal Antibody.



Confocal immunofluorescence analysis of BCBL-1 cells using CD80 Monoclonal Antibody (green), showing membrane localization. Blue: DRAQ5 fluorescent DNA dye.