


Cerberus Monoclonal Antibody

Catalog No :	YM0149
Reactivity :	Human
Applications :	WB;IHC;IF;ELISA
Target :	Cerberus
Fields :	>>Wnt signaling pathway
Gene Name :	CER1
Protein Name :	Cerberus
Human Gene Id :	9350
Human Swiss Prot	O95813
No : Mouse Swiss Prot	O55233
No : Immunogen :	Purified recombinant fragment of human Cerberus expressed in E. Coli.
Specificity :	Cerberus Monoclonal Antibody detects endogenous levels of Cerberus protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Monoclonal, Mouse
Dilution :	WB 1:500 - 1:2000. IHC 1:200 - 1:1000. ELISA: 1:10000 IF 1:50-200
Purification :	Affinity purification
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Molecularweight :	30kD
Cell Pathway :	WNT;WNT-T CELL
· · · · · · · · · · · · · · · · · · ·	




P References :	1. Dev Biol. 1998 Feb 15;194(2):135-51. 2. Growth Factors. 2004 Dec;22(4):233-41. 3. PLoS One. 2009;4(4):e5302.
Background :	This gene encodes a cytokine member of the cysteine knot superfamily, characterized by nine conserved cysteines and a cysteine knot region. The cerberus-related cytokines, together with Dan and DRM/Gremlin, represent a group of bone morphogenetic protein (BMP) antagonists that can bind directly to BMPs and inhibit their activity. [provided by RefSeq, Jul 2008],
Function :	function:Cytokine that may play a role in anterior neural induction and somite formation during embryogenesis in part through a BMP-inhibitory mechanism. Can regulate Nodal signaling during gastrulation as well as the formation and patterning of the primitive streak.,PTM:N-glycosylated.,sequence caution:Translated as Ser.,similarity:Belongs to the DAN family.,similarity:Contains 1 CTCK (C-terminal cystine knot-like) domain.,subunit:Forms monomers and predominantly dimers.,
Subcellular Location :	Secreted .
Expression :	Blood,PCR rescued clones,
Sort :	3862
No4 :	1



Products Images

Western Blot analysis using Cerberus Monoclonal Antibody against CER1 (aa18-267)-hlgGFc transfected HEK293 cell lysate (1).





Immunohistochemistry analysis of paraffin-embedded human gastric cancer (left) and normal gastric tissues (right) with DAB staining using Cerberus Monoclonal Antibody.