

EphA1 Monoclonal Antibody

Catalog No: YM0220

Reactivity: Human

Applications: WB;IHC;IF;ELISA

Target: EphA1

Fields: >>Axon guidance

Gene Name: EPHA1

Protein Name: Ephrin type-A receptor 1

P21709

Q60750

Human Gene Id: 2041

Human Swiss Prot

Iuman Swiss Froi

No:

Mouse Swiss Prot

No:

Immunogen: Purified recombinant fragment of EphA1 expressed in E. Coli.

Specificity: EphA1 Monoclonal Antibody detects endogenous levels of EphA1 protein.

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Monoclonal, Mouse

Dilution: WB 1:500 - 1:2000. IHC 1:200 - 1:1000. ELISA: 1:10000.. IF 1:50-200

Purification : Affinity purification

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight: 108kD

1/3

Cell Pathway: Axon guidance;

P References : 1. Shannon L. Duffy, Kirsten A. Steiner, Patrick P.L. Tam Gene Expr Patterns.

2006 Feb 6.

2. Elena B. Pasquale. Nat Rev Mol Cell Biol.2005 Jun; 6(6): 462-75.

Background: This gene belongs to the ephrin receptor subfamily of the protein-tyrosine kinase

family. EPH and EPH-related receptors have been implicated in mediating developmental events, particularly in the nervous system. Receptors in the EPH subfamily typically have a single kinase domain and an extracellular region containing a Cys-rich domain and 2 fibronectin type III repeats. The ephrin receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. This gene is expressed in some human cancer cell lines and has been implicated

in carcinogenesis. [provided by RefSeq, Jul 2008],

Function: catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine

phosphate.,function:Receptor for members of the ephrin-A family. Binds with a low affinity to ephrin-A1.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. Ephrin receptor subfamily.,similarity:Contains 1 protein

kinase domain.,similarity:Contains 1 SAM (sterile alpha motif) domain.,similarity:Contains 2 fibronectin type-III domains.,tissue

specificity:Overexpressed in several carcinomas.,

Subcellular Location:

Cell membrane; Single-pass type I membrane protein.


Expression : Overexpressed in several carcinomas.

Sort: 5607


No4:

Products Images

2/3



Western Blot analysis using EphA1 Monoclonal Antibody against A549 (1) and HeLa (2) cell lysate.



Immunohistochemistry analysis of paraffin-embedded human ovary carcinoma (left) and breast carcinoma (right), showing cytoplasmic localization with DAB staining using EphA1 Monoclonal Antibody.