

Oct-3/4 Monoclonal Antibody

Catalog No :	YM0485
Reactivity :	Human
Applications :	WB;IF;FCM;ELISA
Target :	Oct-3/4
Fields :	>>Signaling pathways regulating pluripotency of stem cells
Gene Name :	POU5F1
Protein Name :	POU domain class 5 transcription factor 1
Human Gene Id :	5460
Human Swiss Prot No :	Q01860
Mouse Swiss Prot No :	P20263
Immunogen :	Synthesized peptide derived from internal of human Oct-3/4.
Specificity :	Oct-3/4 Monoclonal Antibody detects endogenous levels of Oct-3/4 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Monoclonal, Mouse
Dilution :	WB 1:500 - 1:2000. IF 1:200 - 1:1000. Flow cytometry: 1:200 - 1:400. ELISA: 1:10000. Not yet tested in other applications.
Purification :	Affinity purification
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Molecularweight :	39kD

P References :

1. Stem Cells. 2010 May;28(5):885-93.
2. Mol Med. 2010 Jul-Aug;16(7-8):247-53.
3. Med Sci (Paris). 2010 Apr;26(4):411-6.

Background :

This gene encodes a transcription factor containing a POU homeodomain that plays a key role in embryonic development and stem cell pluripotency. Aberrant expression of this gene in adult tissues is associated with tumorigenesis. This gene can participate in a translocation with the Ewing's sarcoma gene on chromosome 21, which also leads to tumor formation. Alternative splicing, as well as usage of alternative AUG and non-AUG translation initiation codons, results in multiple isoforms. One of the AUG start codons is polymorphic in human populations. Related pseudogenes have been identified on chromosomes 1, 3, 8, 10, and 12. [provided by RefSeq, Oct 2013],

Function :

function:Transcription factor that binds to the octamer motif (5'-ATTTGCAT-3'). Forms a trimeric complex with SOX2 on DNA and controls the expression of a number of genes involved in embryonic development such as YES1, FGF4, UTF1 and ZFP206. Critical for early embryogenesis and for embryonic stem cell pluripotency.,miscellaneous:Several pseudogenes of POU5F1 have been described on chromosomes 1, 3, 8, 10 and 12. 2 of them, localized in chromosomes 8 and 10, are transcribed in cancer tissues but not in normal ones and may be involved in the regulation of POU5F1 gene activity in carcinogenesis.,online information:Oct-4 entry,PTM:Sumoylation enhances the protein stability, DNA binding and transactivation activity. Sumoylation is required for enhanced YES1 expression.,similarity:Belongs to the POU transcription factor family. Class-5 subfamily.,similarity:Contains 1 homeobox DNA-binding doma

Subcellular Location :

Cytoplasm. Nucleus. Expressed in a diffuse and slightly punctuate pattern. Colocalizes with MAPK8 and MAPK9 in the nucleus. .

Expression :

Expressed in developing brain. Highest levels found in specific cell layers of the cortex, the olfactory bulb, the hippocampus and the cerebellum. Low levels of expression in adult tissues.

Tag :

hot

Sort :

11054

No4 :

1


Host :

Mouse


Modifications :

Unmodified


Products Images



Western Blot analysis using Oct-3/4 Monoclonal Antibody against NTERA-2 (1) cell lysate.



Immunofluorescence analysis of NTERA-2 cells using Oct-3/4 Monoclonal Antibody (green). Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.



Flow cytometric analysis of Jurkat cells using Oct-3/4 Monoclonal Antibody (green) and negative control (purple).