

Msi1 Monoclonal Antibody

Catalog No: YM0451

Reactivity: Human

Applications: WB;FCM;ELISA

Target: Msi1

Fields: >>mRNA surveillance pathway

Gene Name: MSI1

Protein Name: RNA-binding protein Musashi homolog 1

O43347

Q61474

Human Gene Id: 4440

Human Swiss Prot

No:

Mouse Swiss Prot

No:

Immunogen: Purified recombinant fragment of human Msi1 expressed in E. Coli.

Specificity: Msi1 Monoclonal Antibody detects endogenous levels of Msi1 protein.

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Monoclonal, Mouse

Dilution: WB 1:500 - 1:2000. Flow cytometry: 1:200 - 1:400. ELISA: 1:10000. Not yet

tested in other applications.

Purification: Affinity purification

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight: 39kD

1/3

P References:

1. BMC Cancer. 2008 Apr 18;8:108.

2. Gastroenterology. 2008 May;134(5):1448-58.

Background:

This gene encodes a protein containing two conserved tandem RNA recognition motifs. Similar proteins in other species function as RNA-binding proteins and play central roles in posttranscriptional gene regulation. Expression of this gene has been correlated with the grade of the malignancy and proliferative activity in gliomas and melanomas. A pseudogene for this gene is located on chromosome 11q13. [provided by RefSeq, Jul 2008],

Function:

domain: The first RNA recognition motif binds more strongly to RNA compared to the second one., function: RNA binding protein that regulates the expression of target mRNAs at the translation level. Regulates expression of the NOTCH1 antagonist NUMB. Binds RNA containing the sequence 5'-GUUAGUUAGUUAGUU-3' and other sequences containing the pattern 5'-[GA]U(1-3)AGU-3'. May play a role in the proliferation and maintenance of stem cells in the central nervous system., similarity: Belongs to the Musashi family., similarity: Contains 2 RRM (RNA recognition motif) domains., tissue specificity: Detected in fetal kidney, brain, liver and lung, and in adult brain and pancreas. Detected in hepatoma cell lines.,

Subcellular Location:

Cytoplasm . Nucleus .

Expression:


Detected in fetal kidney, brain, liver and lung, and in adult brain and pancreas. Detected in hepatoma cell lines.

10285


No4:

Sort:

Products Images



Western Blot analysis using Msi1 Monoclonal Antibody against NTERA-2 cell lysate.



Flow cytometric analysis of PC-2 cells using Msi1 Monoclonal Antibody (green) and negative control (purple).

