

EP-CAM Monoclonal Antibody

Catalog No: YM1034

Reactivity: Human; Mouse

Applications: WB

Target: Ep-CAM

Gene Name: EPCAM

Protein Name: Epithelial cell adhesion molecule

P16422

Human Gene Id: 4072

Human Swiss Prot

No:

Mouse Gene Id: 17075

Mouse Swiss Prot

No:

Immunogen: Purified recombinant human EP-CAM (N-terminus) protein fragments expressed

in E.coli.

Q99JW5

Specificity: EP-CAM Monoclonal Antibody detects endogenous levels of EP-CAM protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Monoclonal, Mouse

Dilution: WB 1:1000 - 1:2000. Not yet tested in other applications.

Purification: Affinity purification

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

1/2

Molecularweight:

35kD

Background:

This gene encodes a carcinoma-associated antigen and is a member of a family that includes at least two type I membrane proteins. This antigen is expressed on most normal epithelial cells and gastrointestinal carcinomas and functions as a homotypic calcium-independent cell adhesion molecule. The antigen is being used as a target for immunotherapy treatment of human carcinomas. Mutations in this gene result in congenital tufting enteropathy. [provided by RefSeq, Dec 2008],

Subcellular Location:

Lateral cell membrane ; Single-pass type I membrane protein . Cell junction, tight junction . Colocalizes with CLDN7 at the lateral cell membrane and tight junction. .

Expression:

Highly and selectively expressed by undifferentiated rather than differentiated embryonic stem cells (ESC). Levels rapidly diminish as soon as ESC's differentiate (at protein levels). Expressed in almost all epithelial cell membranes but not on mesodermal or neural cell membranes. Found on the surface of adenocarcinoma.


Sort:

5593

No4:

1

Products Images



Western Blot analysis using EP-CAM Monoclonal Antibody against HCT116, MCF7 cell lysate.