


UHRF1 Monoclonal Antibody


Catalog No :	YM1111
Reactivity :	Human
Applications :	WB;IF
Target :	UHRF1
Gene Name :	UHRF1
Protein Name :	E3 ubiquitin-protein ligase UHRF1
Human Gene Id :	29128
Human Swiss Prot No :	Q96T88
Mouse Swiss Prot No :	Q8VDF2
Immunogen :	Purified recombinant human UHRF1 (N-terminus) protein fragments expressed in E.coli.
Specificity :	UHRF1 Monoclonal Antibody detects endogenous levels of UHRF1 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Monoclonal, Mouse
Dilution :	WB 1:1000 - 1:2000. IF 1:100 - 1:500. Not yet tested in other applications.
Purification :	Affinity purification
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Molecularweight :	90kD

Background :	This gene encodes a member of a subfamily of RING-finger type E3 ubiquitin ligases. The protein binds to specific DNA sequences, and recruits a histone deacetylase to regulate gene expression. Its expression peaks at late G1 phase and continues during G2 and M phases of the cell cycle. It plays a major role in the G1/S transition by regulating topoisomerase IIalpha and retinoblastoma gene expression, and functions in the p53-dependent DNA damage checkpoint. It is regarded as a hub protein for the integration of epigenetic information. This gene is up-regulated in various cancers, and it is therefore considered to be a therapeutic target. Multiple transcript variants encoding different isoforms have been found for this gene. A related pseudogene exists on chromosome 12. [provided by RefSeq, Feb 2014],
Function :	developmental stage:Expressed in fetal thymus, liver and kidney.,domain:The RING finger is required for ubiquitin ligase activity.,domain:The YDG domain mediates the interaction with histone H3.,function:Putative E3 ubiquitin-protein ligase. May participate in methylation-dependent transcriptional regulation. Binds to inverted 5'-CCAAT-3' box 2 in the TOP2A promoter, and activates TOP2A expression. Important for G1/S transition. May be involved in DNA repair and chromosomal stability.,induction:Up-regulated in proliferating cells, and down-regulated in quiescent cells. Down-regulated upon adriamycin-induced DNA damage, in a TP53/p53 and CDKN1A-dependent way. Induced by E2F1 transcription factor.,pathway:Protein modification; protein ubiquitination.,PTM:Phosphorylated on serine residues. Phosphorylation may enhance DNA-binding activity.,PTM:Ubiquitinated; which leads to proteasomal degrad
Subcellular Location :	Nucleus . Localizes to replication foci. Enriched in pericentric heterochromatin. Also localizes to euchromatic regions.
Expression :	Expressed in thymus, bone marrow, testis, lung and heart. Overexpressed in breast cancer.
Tag :	orthogonal
Sort :	1259
No4 :	1

Products Images



Western Blot analysis using UHRF1 Monoclonal Antibody against mouse heart, 293T, HeLa cell lysate, HeLa nuclear extract.



Immunofluorescence analysis of HeLa cells using UHRF1 Monoclonal Antibody.