

## COX IV Monoclonal Antibody(6C8), Biotin Conjugated

Catalog No: YM2011

**Reactivity:** Human;Rat;Mouse

**Applications:** WB;IHC;IF;

Target: COX IV

**Fields:** >>Oxidative phosphorylation;>>Metabolic pathways;>>Cardiac muscle

contraction;>>Thermogenesis;>>Non-alcoholic fatty liver disease;>>Alzheimer disease;>>Parkinson disease;>>Amyotrophic lateral sclerosis;>>Huntington disease;>>Prion disease;>>Pathways of neurodegeneration - multiple diseases;>>Chemical carcinogenesis - reactive oxygen species;>>Diabetic

cardiomyopathy

Gene Name: COX4I1

**Protein Name:** Cytochrome c oxidase subunit 4 isoform 1, mitochondrial

Human Gene Id: 1327

**Human Swiss Prot** 

No:

**Specificity:** COX IV Monoclonal Antibody(6C8) Biotin conjugated specially designed for your

WB or IHC analysis.

**Formulation:** Liquid in PBS, pH 7.4, containing 0.02% sodium azide as preservative and 50%

Glycerol.

P13073

Source: Monoclonal, Mouse IgG1

**Dilution:** Optimal working dilutions should be determined experimentally by the

investigator. Suggested starting dilutions are as follows: WB 1:1000-3000, IHC

1:50-300.

**Purification:** The antibody was affinity-purified from mouse ascites by affinity-

chromatography using specific immunogen.

**Concentration**: 1mg/ml

1/2



**Storage Stability:** 

Stable for one year at -15°C to -25°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezi

**Cell Pathway:** 

Oxidative phosphorylation; Cardiac muscle contraction; Alzheimer's disease; Parkinson's disease; Huntington's disease;

**Background:** 

Cytochrome c oxidase (COX) is the terminal enzyme of the mitochondrial respiratory chain. It is a multi-subunit enzyme complex that couples the transfer of electrons from cytochrome c to molecular oxygen and contributes to a proton electrochemical gradient across the inner mitochondrial membrane. The complex consists of 13 mitochondrial- and nuclear-encoded subunits. The mitochondrially-encoded subunits perform the electron transfer and proton pumping activities. The functions of the nuclear-encoded subunits are unknown but they may play a role in the regulation and assembly of the complex. This gene encodes the nuclear-encoded subunit IV isoform 1 of the human mitochondrial respiratory chain enzyme. It is located at the 3' of the NOC4 (neighbor of COX4) gene in a head-to-head orientation, and shares a promoter with it. Pseudogenes related to this gene are located on chromosomes

**Function:** 

function: This protein is one of the nuclear-coded polypeptide chains of cytochrome c oxidase, the terminal oxidase in mitochondrial electron transport., similarity: Belongs to the cytochrome c oxidase IV family., tissue specificity: Ubiquitous.,

Subcellular Location:

Sort:

Mitochondrion inner membrane; Single-pass membrane protein.

Expression:

4462

Ubiquitous.

No4:

## **Products Images**

2/2