


## OR5B2 Polyclonal Antibody

<b>Catalog No :</b>	YN2643
<b>Reactivity :</b>	Human;Rat;Mouse;
<b>Applications :</b>	WB;ELISA
<b>Target :</b>	OR5B2
<b>Fields :</b>	>>Olfactory transduction
<b>Gene Name :</b>	OR5B2
<b>Protein Name :</b>	Olfactory receptor 5B2 (OST073) (Olfactory receptor OR11-240)
<b>Human Gene Id :</b>	390190
<b>Human Swiss Prot No :</b>	Q96R09
<b>Immunogen :</b>	Synthesized peptide derived from human protein . at AA range: 130-210
<b>Specificity :</b>	OR5B2 Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Observed Band :</b>	33kD

<b>Background :</b>	olfactory receptor family 5 subfamily B member 2(OR5B2) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],
<b>Function :</b>	function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,
<b>Subcellular Location :</b>	Cell membrane; Multi-pass membrane protein.
<b>Sort :</b>	20670
<b>No4 :</b>	1

## Products Images



Western blot analysis of lysates from PC12 cells, primary antibody was diluted at 1:1000, 4° over night