

## NARG2 rabbit pAb

|                              |   |
|------------------------------|---|
| <b>Catalog No :</b>          | YN5286  |
| <b>Reactivity :</b>          | Human;Mouse   |
| <b>Applications :</b>        | WB  |
| <b>Target :</b>              | NARG2   |
| <b>Gene Name :</b>           | NARG2 BRCC1 UNQ3101/PRO10100  |
| <b>Protein Name :</b>        | NMDA receptor-regulated protein 2   |
| <b>Human Gene Id :</b>       | 79664   |
| <b>Human Swiss Prot No :</b> | Q659A1  |
| <b>Mouse Gene Id :</b>       | 93697   |
| <b>Mouse Swiss Prot No :</b> | Q3UZ18  |
| <b>Rat Gene Id :</b>         | 93697   |
| <b>Immunogen :</b>           | Synthesized peptide derived from human NARG2 AA range: 474-524  |
| <b>Specificity :</b>         | This antibody detects endogenous levels of human NARG2  |
| <b>Formulation :</b>         | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| <b>Source :</b>              | Polyclonal, Rabbit,IgG  |
| <b>Dilution :</b>            | WB 1:1000-2000  |
| <b>Purification :</b>        | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| <b>Concentration :</b>       | 1 mg/ml   |

**Storage Stability :** -15°C to -25°C/1 year(Do not lower than -25°C)

**Molecularweight :** 108kD

**Subcellular Location :** Nucleus . Colocalizes with COIL in subnuclear Cajal and histone locus bodies. Translocates in the LEC complex to Cajal and histone locus bodies at snRNA genes in a ICE1-dependent manner. Associates to transcriptionally active chromatin at snRNA genes.

**Expression :** Expressed at low levels in lung and testis.

**Sort :** 17636

**No4 :** 1

## Products Images