


Oct1 Monoclonal Antibody(7G1)


Catalog No :	YM3089
Reactivity :	Human
Applications :	WB;IHC;IF;
Target :	OCT1
Fields :	>>Herpes simplex virus 1 infection;>>Lipid and atherosclerosis
Gene Name :	POU2F1
Protein Name :	POU domain class 2 transcription factor
Human Gene Id :	5451/5452
Human Swiss Prot No :	P14859/P09086
Mouse Gene Id :	18986/18987
Rat Gene Id :	171068
Rat Swiss Prot No :	P31503
Immunogen :	Synthetic Peptide of 42644
Specificity :	The antibody detects endogenous Oct1 protein.
Formulation :	PBS, pH 7.4, containing 0.5%BSA, 0.02% sodium azide as Preservative and 50% Glycerol.
Source :	Monoclonal, Mouse
Dilution :	WB 1:500-2000 IF 1:200 IHC 1:50-300
Purification :	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.

Storage Stability :	-15 °C to -25 °C/1 year(Do not lower than -25 °C)
Observed Band :	89kD
Background :	The OCT1 transcription factor was among the first identified members of the POU transcription factor family (summarized by Sturm et al., 1993 [PubMed 8314572]). Members of this family contain the POU domain, a 160-amino acid region necessary for DNA binding to the octameric sequence ATGCAAAT.[supplied by OMIM, Jul 2010],
Function :	function:Transcription factor that binds to the octamer motif (5'-ATTTGCAT-3') and activates the promoters of the genes for some small nuclear RNAs (snRNA) and of genes such as those for histone H2B and immunoglobulins. Modulates transcription transactivation by NR3C1, AR and PGR.,PTM:Phosphorylated by PRKDC.,similarity:Belongs to the POU transcription factor family. Class-2 subfamily.,similarity:Contains 1 homeobox DNA-binding domain.,similarity:Contains 1 POU-specific domain.,subunit:Interacts with NR3C1, AR, PGR and HCFC1.,tissue specificity:Ubiquitous. Isoform 2 is lymphocyte-specific.,
Subcellular Location :	Nucleus.
Expression :	Ubiquitous. Isoform 2 is lymphocyte-specific.
Tag :	hot
Sort :	11044
Host :	Mouse
Modifications :	Unmodified


Products Images



Immunohistochemical analysis of paraffin-embedded Human-breast-cancer tissue. 1, Oct1 Monoclonal Antibody(7G1) was diluted at 1:200(4°C, overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C, 20min). 3, Secondary antibody was diluted at 1:200(room temperature, 30min). Negative control was used by secondary antibody only.



Immunofluorescence analysis of Human-breast-cancer tissue. 1, Oct1 Monoclonal Antibody(7G1)(red) was diluted at 1:200(4°C, overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min). 3, Picture B: DAPI(blue) 10min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B



Western blot analysis of 1) HeLa, 2) Jurkat, 3) HepG2, diluted at 1:2000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Invent biotech, MN, USA).