

PHF10 rabbit pAb

Catalog No :	YN8733
Reactivity :	Human;Mouse;Rat
Applications :	WB
Target :	PHF10
Gene Name :	PHF10 BAF45A
Protein Name :	PHD finger protein 10 (BRG1-associated factor 45a) (BAF45a) (XAP135)
Human Gene Id :	55274
Human Swiss Prot	Q8WUB8
No : Mouse Gene Id :	72057
Mouse Swiss Prot	Q9D8M7
No : Rat Gene Id :	292404
Rat Swiss Prot No :	Q4V7A6
Immunogen :	Synthesized peptide derived from human PHF10
Specificity :	This antibody detects endogenous levels of PHF10 at Human, Mouse, Rat
Formulation :	Liquid in PBS containing 50% glycerol, 0,5% BSA and 0.02% sodium azide.
Source :	Polyclonal. Rabbit.loG
Dilution :	WB 1:500-2000
Duvifientiers	
Purification :	chromatography using epitope-specific immunogen.



Concentration : 1 mg/ml Storage Stability : -15°C to -25°C/1 year(Do not lower than -25°C) Molecularweight : 55kD Function : Involved in transcription activity regulation by chromatin remodeling. Belongs the neural progenitors-specific chromatin remodeling complex (npBAF compleand is required for the proliferation of neural progenitors. During neural development a switch from a stem/progenitor to a post-mitotic chromatin remodeling mechanism occurs as neurons exit the cell cycle and become	; to ;x)
Storage Stability : -15°C to -25°C/1 year(Do not lower than -25°C) Molecularweight : 55kD Function : Involved in transcription activity regulation by chromatin remodeling. Belongs the neural progenitors-specific chromatin remodeling complex (npBAF comple and is required for the proliferation of neural progenitors. During neural development a switch from a stem/progenitor to a post-mitotic chromatin remodeling mechanism occurs as neurons exit the cell cycle and become	; to ;x)
Storage Stability : -15°C to -25°C/1 year(Do not lower than -25°C) Molecularweight : 55kD Function : Involved in transcription activity regulation by chromatin remodeling. Belongs the neural progenitors-specific chromatin remodeling complex (npBAF complex and is required for the proliferation of neural progenitors. During neural development a switch from a stem/progenitor to a post-mitotic chromatin remodeling mechanism occurs as neurons exit the cell cycle and become	s to ex)
Molecularweight : 55kD Function : Involved in transcription activity regulation by chromatin remodeling. Belongs the neural progenitors-specific chromatin remodeling complex (npBAF complex and is required for the proliferation of neural progenitors. During neural development a switch from a stem/progenitor to a post-mitotic chromatin remodeling mechanism occurs as neurons exit the cell cycle and become	; to ;x)
Molecularweight : 55kD Function : Involved in transcription activity regulation by chromatin remodeling. Belongs the neural progenitors-specific chromatin remodeling complex (npBAF comple and is required for the proliferation of neural progenitors. During neural development a switch from a stem/progenitor to a post-mitotic chromatin remodeling mechanism occurs as neurons exit the cell cycle and become	; to ;x)
Function : Involved in transcription activity regulation by chromatin remodeling. Belongs the neural progenitors-specific chromatin remodeling complex (npBAF complex and is required for the proliferation of neural progenitors. During neural development a switch from a stem/progenitor to a post-mitotic chromatin remodeling mechanism occurs as neurons exit the cell cycle and become	s to ≆x)
Function : Involved in transcription activity regulation by chromatin remodeling. Belongs the neural progenitors-specific chromatin remodeling complex (npBAF complex and is required for the proliferation of neural progenitors. During neural development a switch from a stem/progenitor to a post-mitotic chromatin remodeling mechanism occurs as neurons exit the cell cycle and become	s to ∋x)
committed to their adult state. The transition from proliferating neural stem/progenitor cells to post-mitotic neurons requires a switch in subunit composition of the npBAF and nBAF complexes. As neural progenitors exit mitosis and differentiate into neurons, npBAF complexes which contain ACTL6A/BAF53A and PHF10/BAF45A, are exchanged for homologous alternative ACTL6B/BAF53B and DPF1/BAF45B or DPF3/BAF45C subunits neuron-specific complexes (nBAF). The npBAF complex is essential for the se renewal/proliferative cap	in əlf-
Subcellular Nucleus .	
Location :	
Sort : 27410	
No4 : 1	

Products Images