

PAK6 Polyclonal Antibody

Catalog No :	YN0034
Reactivity :	Human;Mouse
Applications :	WB;ELISA
Target :	PAK6
Fields :	>>ErbB signaling pathway;>>Ras signaling pathway;>>Axon guidance;>>Focal adhesion;>>T cell receptor signaling pathway;>>Regulation of actin cytoskeleton;>>Human immunodeficiency virus 1 infection;>>Renal cell carcinoma
Gene Name :	PAK6 PAK5
Protein Name :	Serine/threonine-protein kinase PAK 6 (EC 2.7.11.1) (PAK-5) (p21-activated kinase 6) (PAK-6)
Human Gene Id :	56924
Human Swiss Prot No :	Q9NQU5
Mouse Swiss Prot No :	Q3ULB5
Immunogen :	Synthesized peptide derived from human protein . at AA range: 100-180
Specificity :	PAK6 Polyclonal Antibody detects endogenous levels of protein.
Formulation :	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000 ELISA 1:5000-20000
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml

Storage Stability : -15°C to -25°C/1 year (Do not lower than -25°C)

Observed Band : 74kD

Cell Pathway : ErbB_HER; Axon guidance; Focal adhesion; T_Cell_Receptor; Regulates Actin and Cytoskeleton; Renal cell carcinoma;

Background : This gene encodes a member of a family of p21-stimulated serine/threonine protein kinases, which contain an amino-terminal Cdc42/Rac interactive binding (CRIB) domain and a carboxyl-terminal kinase domain. These kinases function in a number of cellular processes, including cytoskeleton rearrangement, apoptosis, and the mitogen-activated protein (MAP) kinase signaling pathway. The protein encoded by this gene interacts with androgen receptor (AR) and translocates to the nucleus, where it is involved in transcriptional regulation. Changes in expression of this gene have been linked to prostate cancer. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Dec 2015],

Function : catalytic activity: ATP + a protein = ADP + a phosphoprotein., function: The activated kinase acts on a variety of targets., PTM: Autophosphorylated when activated by CDC42/p21., similarity: Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. STE20 subfamily., similarity: Contains 1 CRIB domain., similarity: Contains 1 protein kinase domain., subunit: Interacts tightly with GTP-bound but not GDP-bound CDC42/p21 and RAC1 (By similarity). Interacts with the androgen receptor.,

Subcellular Location : Cytoplasm. Nucleus. Cotranslocates into nucleus with AR in response to androgen induction.

Expression : Selectively expressed in brain and testis, with lower levels in multiple tissues including prostate and breast.

Products Images

