


PPAR-α Polyclonal Antibody


Catalog No :	YT3835
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC;IF;ELISA
Target :	PPAR α
Fields :	>>PPAR signaling pathway;>>cAMP signaling pathway;>>Adipocytokine signaling pathway;>>Glucagon signaling pathway;>>Insulin resistance;>>Non-alcoholic fatty liver disease;>>Alcoholic liver disease;>>Hepatitis C;>>Chemical carcinogenesis - receptor activation;>>Diabetic cardiomyopathy
Gene Name :	PPARA
Protein Name :	Peroxisome proliferator-activated receptor alpha
Human Gene Id :	5465
Human Swiss Prot No :	Q07869
Mouse Gene Id :	19013
Mouse Swiss Prot No :	P23204
Rat Gene Id :	25747
Rat Swiss Prot No :	P37230
Immunogen :	The antiserum was produced against synthesized peptide derived from human PPAR-alpha. AA range:6-55
Specificity :	PPAR-α Polyclonal Antibody detects endogenous levels of PPAR-α protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG

Dilution :	WB 1:500-2000, ELISA 1:10000-20000 IHC 1:50-300. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15 °C to -25 °C/1 year(Do not lower than -25 °C)
Observed Band :	52kD
Cell Pathway :	PPAR;Adipocytokine;
Background :	<p>peroxisome proliferator activated receptor alpha(PPARA) Homo sapiens Peroxisome proliferators include hypolipidemic drugs, herbicides, leukotriene antagonists, and plasticizers; this term arises because they induce an increase in the size and number of peroxisomes. Peroxisomes are subcellular organelles found in plants and animals that contain enzymes for respiration and for cholesterol and lipid metabolism. The action of peroxisome proliferators is thought to be mediated via specific receptors, called PPARs, which belong to the steroid hormone receptor superfamily. PPARs affect the expression of target genes involved in cell proliferation, cell differentiation and in immune and inflammation responses. Three closely related subtypes (alpha, beta/delta, and gamma) have been identified. This gene encodes the subtype PPAR-alpha, which is a nuclear transcription factor. Multiple alternatively spliced transcript variants have been described for thi</p>
Function :	<p>function:Receptor that binds peroxisome proliferators such as hypolipidemic drugs and fatty acids. Once activated by a ligand, the receptor binds to a promoter element in the gene for acyl-CoA oxidase and activates its transcription. It therefore controls the peroxisomal beta-oxidation pathway of fatty acids.,online information:Peroxisome proliferator-activated receptor entry,similarity:Belongs to the nuclear hormone receptor family. NR1 subfamily.,similarity:Contains 1 nuclear receptor DNA-binding domain.,subunit:Heterodimer with the retinoid X receptor. Interacts with NCOA3 and NCOA6 coactivators, leading to a strong increase of transcription of target genes. Also interacts with PPARBP coactivator in vitro. Interacts with AKAP13.,tissue specificity:Skeletal muscle, liver, heart and kidney.,</p>
Subcellular Location :	Nucleus.
Expression :	Skeletal muscle, liver, heart and kidney. Expressed in monocytes (PubMed:28167758).


Products Images




Immunofluorescence analysis of HeLa cell. 1,PPAR- α Polyclonal Antibody(red) was diluted at 1:200(4 ° overnight). Galectin-3 Monoclonal Antibody(6G2)(green) was diluted at 1:200(4 ° overnight). 2, Goat Anti Rabbit Alexa Fluor 594 Catalog:RS3611 was diluted at 1:1000(room temperature, 50min). Goat Anti Mouse Alexa Fluor 488 Catalog:RS3208 was diluted at 1:1000(room temperature, 50min).




Immunofluorescence analysis of rat-spleen tissue. 1,PPAR- α Polyclonal Antibody(red) was diluted at 1:200(4 ° C,overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min).3, Picture B: DAPI(blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B




Immunohistochemical analysis of paraffin-embedded Mouse-kidney tissue. 1,PPAR- α Polyclonal Antibody was diluted at 1:200(4 ° C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98 ° C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min). Negative control was used by secondary antibody only.



Western Blot analysis of 3T3 cells using PPAR-α Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).



Immunofluorescence analysis of HeLa cells, using PPAR-alpha Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from NIH/3T3 cells, using PPAR-alpha Antibody. The lane on the right is blocked with the synthesized peptide.