

Ribosomal Protein L7 Polyclonal Antibody

Catalog No :	YT4118
Reactivity :	Human;Mouse;Rat;Cat
Applications :	WB;IHC;IF;ELISA
Target :	Ribosomal Protein L7
Fields :	>>Ribosome;>>Coronavirus disease - COVID-19
Gene Name :	RPL7
Protein Name :	60S ribosomal protein L7
Human Gene Id :	6129
Human Swiss Prot	P18124
No : Mouse Gene Id :	19989
Mouse Swiss Prot	P14148
No : Rat Swiss Prot No :	P05426
Immunogen :	The antiserum was produced against synthesized peptide derived from human RPL7. AA range:199-248
Specificity :	Ribosomal Protein L7 Polyclonal Antibody detects endogenous levels of Ribosomal Protein L7 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:10000 IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity- chromatography using epitope-specific immunogen.



Best Tools for immunology Research		
Concentration :	1 mg/ml	
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)	
Observed Band :	32kD	
Coll Pathway :	Bibosome:	
Cen Fathway.		
Background :	Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L30P family of ribosomal proteins. It contains an N-terminal basic region-leucine zipper (BZIP)-like domain and the RNP consensus submotif RNP2. In vitro the BZIP-like domain mediates homodimerization and stable binding to DNA and RNA, with a preference for 28S rRNA and mRNA. The protein can inhibit cell-free translation of mRNAs, suggesting that it plays a regulatory role in the translation apparatus. It is located in the cytoplasm. The protein has been shown to be an autoantigen in patients with systemic autoimmune diseases, such as systemic lupus erythematosus. As is typical	
Function :	function:Binds to G-rich structures in 28S rRNA and in mRNAs. Plays a regulatory role in the translation apparatus; inhibits cell-free translation of mRNAs.,similarity:Belongs to the ribosomal protein L30P family.,subunit:Homodimer.,	
Subcellular Location :	nucleus,nucleolus,cytoplasm,cytosol,ribosome,focal adhesion,membrane,cytosolic large ribosomal subunit,intracellular ribonucleoprotein complex,extracellular exosome,	
Expression :	Bone marrow,Eye,Fibroblast,Hepatoma,Lung,Mammary gland,Muscle,Pancreas,Skin,	

Products Images







25---

15-

15---

Western blot analysis of Mouse-kidney mouse-brain HELA KB SH-SY5Y 293T 3T3 lysis using Ribosomal Protein L7 antibody. Antibody was diluted at 1:2000

Ribosomal Protein L7





Western blot analysis of lysates from 293 and HeLa cells, using RPL7 Antibody. The lane on the right is blocked with the synthesized peptide.

Western blot analysis of the lysates from K562 cells using RPL7 antibody.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Tris-EDTA,pH9.0 was used for antigen retrieval. 2 Antibody was diluted at 1:200(4° overnight.3,Secondary antibody was diluted at 1:200(room temperature, 45min).