


## Wee 1 Polyclonal Antibody

<b>Catalog No :</b>	YT4903
<b>Reactivity :</b>	Human;Mouse;Rat
<b>Applications :</b>	WB;ELISA
<b>Target :</b>	WEE1
<b>Fields :</b>	>>Cell cycle;>>Human immunodeficiency virus 1 infection
<b>Gene Name :</b>	WEE1
<b>Protein Name :</b>	Wee1-like protein kinase
<b>Human Gene Id :</b>	7465
<b>Human Swiss Prot No :</b>	P30291
<b>Mouse Gene Id :</b>	22390
<b>Mouse Swiss Prot No :</b>	P47810
<b>Rat Gene Id :</b>	308937
<b>Rat Swiss Prot No :</b>	Q63802
<b>Immunogen :</b>	The antiserum was produced against synthesized peptide derived from human WEE1. AA range:19-68
<b>Specificity :</b>	Wee 1 Polyclonal Antibody detects endogenous levels of Wee 1 protein.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500 - 1:2000. ELISA: 1:40000. Not yet tested in other applications.

<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Observed Band :</b>	72kD
<b>Cell Pathway :</b>	Cell_Cycle_G1S;Cell_Cycle_G2M_DNA;
<b>Background :</b>	WEE1 G2 checkpoint kinase(WEE1) Homo sapiens This gene encodes a nuclear protein, which is a tyrosine kinase belonging to the Ser/Thr family of protein kinases. This protein catalyzes the inhibitory tyrosine phosphorylation of CDC2/cyclin B kinase, and appears to coordinate the transition between DNA replication and mitosis by protecting the nucleus from cytoplasmically activated CDC2 kinase. [provided by RefSeq, Jul 2008],
<b>Function :</b>	catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate.,cofactor:Binds 2 magnesium ions per subunit.,enzyme regulation:Synthesis is increased during S and G2 phases, presumably by an increase in transcription; activity is decreased by phosphorylation during m phase. Protein levels fall in M phase as a result of decreased synthesis combined with degradation. Activity seems to be negatively regulated by phosphorylation upon entry into mitosis, although N-terminal phosphorylation might also regulate the protein stability via protection from proteolysis or might regulate the subcellular location.,function:May act as a negative regulator of entry into mitosis (G2 to M transition) by protecting the nucleus from cytoplasmically activated cyclin B1-complexed CDC2 before the onset of mitosis. Its activity increases during S and G2 phases and decreases at M phase
<b>Subcellular Location :</b>	Nucleus.
<b>Expression :</b>	Amygdala,Blood,Epithelium,Human uterus endothel primary cell culture,Placenta,Skin,

## Products Images



Western blot analysis of lysates from 293 cells, using WEE1 Antibody. The lane on the right is blocked with the synthesized peptide.