

## FA2H Polyclonal Antibody

Catalog No :	YT6078
Reactivity :	Human;Mouse;Rat
Applications :	WB;ELISA
Target :	FA2H
Gene Name :	FA2H
Protein Name :	FA2H
Human Gene Id :	79152
Human Swiss Prot	Q7L5A8
No : Mouse Gene Id :	338521
Mouse Swiss Prot	Q5MPP0
NO : Immunogen :	Synthesized peptide derived from human FA2H. at AA range: 101-150
Specificity :	FA2H Polyclonal Antibody detects endogenous levels of FA2H
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000, ELISA 1:10000-20000
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity- chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)



<b>Observed Band :</b>	55kD
Background :	This gene encodes a protein that catalyzes the synthesis of 2-hydroxysphingolipids, a subset of sphingolipids that contain 2-hydroxy fatty acids. Sphingolipids play roles in many cellular processes and their structural diversity arises from modification of the hydrophobic ceramide moiety, such as by 2-hydroxylation of the N-acyl chain, and the existence of many different head groups. Mutations in this gene have been associated with leukodystrophy dysmyelinating with spastic paraparesis with or without dystonia.[provided by RefSeq, Mar 2010],
Function :	cofactor:Iron.,disease:Defects in FA2H are the cause of leukodystrophy dysmyelinating with spastic paraparesis with or without dystonia (DLDSP) [MIM:612443]. The disorder consists of a progressive neurologic disease manifested by spasticity, disordered tonicity of muscle, and white matter degeneration.,domain:The histidine box domains may contain the active site and/or be involved in metal ion binding.,function:Required for alpha-hydroxylation of free fatty acids and the formation of alpha-hydroxylated sphingolipids.,induction:Up-regulated during keratinocyte differentiation.,similarity:Belongs to the SCS7 family.,similarity:Contains 1 cytochrome b5 heme-binding domain.,tissue specificity:Detected in differentiating cultured keratinocytes (at protein level). Detected in epidermis and cultured keratinocytes. Highly expressed in brain and colon. Detected at lower levels in testis, prostate
Subcellular Location :	Endoplasmic reticulum membrane ; Multi-pass membrane protein . Microsome membrane ; Multi-pass membrane protein .
Expression :	Detected in differentiating cultured keratinocytes (at protein level). Detected in epidermis and cultured keratinocytes (PubMed:17355976). Highly expressed in brain and colon. Detected at lower levels in testis, prostate, pancreas and kidney (PubMed:15337768).

## **Products Images**

Western Blot analysis of mouse-heart cells using primary antibody diluted at 1:2000(4°C overnight). Secondary antibody:Goat Anti-rabbit IgG IRDye 800( diluted at 1:5000, 25°C, 1 hour)

