

CD14 rabbit pAb

Catalog No :	YT7957
Reactivity :	Human;Rat;Mouse;
Applications :	WB;ELISA
Target :	CD14
Fields :	>>MAPK signaling pathway;>>NF-kappa B signaling pathway;>>Phagosome;>>Toll-like receptor signaling pathway;>>Hematopoietic cell lineage;>>Alcoholic liver disease;>>Shigellosis;>>Salmonella infection;>>Pertussis;>>Legionellosis;>>Amoebiasis;>>Tuberculosis;>>Transcriptional misregulation in cancer;>>Acute myeloid leukemia;>>Lipid and atherosclerosis
Gene Name :	CD14
Protein Name :	CD14
Human Gene Id :	929
Human Swiss Prot No :	P08571
Mouse Gene Id :	12475
Mouse Swiss Prot No :	P10810
Rat Gene Id :	60350
Rat Swiss Prot No :	Q63691
Immunogen :	Synthesized peptide derived from human CD14
Specificity :	This antibody detects endogenous levels of Human CD14
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG

Dilution :	WB 1:1000-2000 ELISA 1:5000-20000
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15 °C to -25 °C/1 year(Do not lower than -25 °C)
Molecularweight :	41kD
Background :	The protein encoded by this gene is a surface antigen that is preferentially expressed on monocytes/macrophages. It cooperates with other proteins to mediate the innate immune response to bacterial lipopolysaccharide. Alternative splicing results in multiple transcript variants encoding the same protein. [provided by RefSeq, Mar 2010],
Function :	function:Cooperates with MD-2 and TLR4 to mediate the innate immune response to bacterial lipopolysaccharide (LPS). Acts via MyD88, TIRAP and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response. Up-regulates cell surface molecules, including adhesion molecules.,online information:CD14 entry,similarity:Contains 11 LRR (leucine-rich) repeats.,subunit:Belongs to the lipopolysaccharide (LPS) receptor, a multi-protein complex containing at least CD14, MD-2 and TLR4.,tissue specificity:Expressed strongly on the surface of monocytes and weakly on the surface of granulocytes; also expressed by most tissue macrophages.,
Subcellular Location :	Cell membrane ; Lipid-anchor, GPI-anchor . Secreted . Membrane raft . Golgi apparatus . Secreted forms may arise by cleavage of the GPI anchor. .
Expression :	Detected on macrophages (at protein level) (PubMed:1698311). Expressed strongly on the surface of monocytes and weakly on the surface of granulocytes; also expressed by most tissue macrophages.

Products Images