

BLNK (phospho Tyr96) Polyclonal Antibody

Catalog No :	YP0687
Reactivity :	Human;Mouse;Monkey
Applications :	WB;IHC;IF;ELISA
Target :	BLNK
Fields :	>>NF-kappa B signaling pathway;>>Osteoclast differentiation;>>B cell receptor signaling pathway;>>Epstein-Barr virus infection;>>Primary immunodeficiency
Gene Name :	BLNK
Protein Name :	B-cell linker protein
Human Gene Id :	29760
Human Swiss Prot	Q8WV28
Mouse Gene Id :	17060
Mouse Swiss Prot	Q9QUN3
NO : Immunogen :	The antiserum was produced against synthesized peptide derived from human BLNK around the phosphorylation site of Tyr96. AA range:62-111
Specificity :	Phospho-BLNK (Y96) Polyclonal Antibody detects endogenous levels of BLNK protein only when phosphorylated at Y96.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:20000 IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity- chromatography using epitope-specific immunogen.



Best Tools for immunology Research		
Concentration :	1 mg/ml	
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)	
Observed Band :	50kD	
Cell Pathway :	B_Cell_Antigen;Primary immunodeficiency;	
Background :	This gene encodes a cytoplasmic linker or adaptor protein that plays a critical role in B cell development. This protein bridges B cell receptor-associated kinase activation with downstream signaling pathways, thereby affecting various biological functions. The phosphorylation of five tyrosine residues is necessary for this protein to nucleate distinct signaling effectors following B cell receptor activation. Mutations in this gene cause hypoglobulinemia and absent B cells, a disease in which the pro- to pre-B-cell transition is developmentally blocked. Deficiency in this protein has also been shown in some cases of pre-B acute lymphoblastic leukemia. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, May 2012],	
Function :	disease:Defects in BLNK are the cause of hypoglobulinemia and absent B-cells [MIM:604515]. This is a developmental blockage at the pro- to pre-B-cell transition.,disease:In 6 of 34 childhood pre-B acute lymphoblastic leukemia (ALL) samples that were tested showed a complete loss or drastic reduction of BLNK expression.,function:Functions as a central linker protein that bridges kinases associated with the B-cell receptor (BCR) with a multitude of signaling pathways, regulating biological outcomes of B-cell function and development. Plays a role in the activation of ERK/EPHB2, MAP kinase p38 and JNK. Modulates AP1 activation. Important for the activation of NF-kappa-B and NFAT. Plays an important role in BCR-mediated PLCG1 and PLCG2 activation and Ca(2+) mobilization and is required for trafficking of the BCR to late endosomes. However, does not seem to be required for pre-BCR-mediated ac	
Subcellular Location :	Cytoplasm . Cell membrane . BCR activation results in the translocation to membrane fraction.	
Expression :	Expressed in B-cell lineage and fibroblast cell lines (at protein level). Highest levels of expression in the spleen, with lower levels in the liver, kidney, pancreas, small intestines and colon.	

Products Images





Western Blot analysis of COLO205 cells using Phospho-BLNK (Y96) Polyclonal Antibody



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using BLNK (Phospho-Tyr96) Antibody



Immunohistochemistry analysis of paraffin-embedded human lymph node, using BLNK (Phospho-Tyr96) Antibody. The picture on the right is blocked with the phospho peptide.





Western blot analysis of lysates from COLO205 cells, using BLNK (Phospho-Tyr96) Antibody. The lane on the right is blocked with the phospho peptide.