


## Na<sup>+</sup>/K<sup>+</sup>-ATPase α1 (phospho Ser16) Polyclonal Antibody

<b>Catalog No :</b>	YP0644
<b>Reactivity :</b>	Human;Mouse;Rat
<b>Applications :</b>	WB;IHC;IF;ELISA
<b>Target :</b>	Na <sup>+</sup> /K <sup>+</sup> -ATPase α1
<b>Fields :</b>	>>cGMP-PKG signaling pathway;>>cAMP signaling pathway;>>Cardiac muscle contraction;>>Adrenergic signaling in cardiomyocytes;>>Insulin secretion;>>Thyroid hormone synthesis;>>Thyroid hormone signaling pathway;>>Aldosterone synthesis and secretion;>>Aldosterone-regulated sodium reabsorption;>>Endocrine and other factor-regulated calcium reabsorption;>>Proximal tubule bicarbonate reclamation;>>Salivary secretion;>>Gastric acid secretion;>>Pancreatic secretion;>>Carbohydrate digestion and absorption;>>Protein digestion and absorption;>>Bile secretion;>>Mineral absorption
<b>Gene Name :</b>	ATP1A1
<b>Protein Name :</b>	Sodium/potassium-transporting ATPase subunit alpha-1
<b>Human Gene Id :</b>	476
<b>Human Swiss Prot No :</b>	P05023
<b>Mouse Gene Id :</b>	11928
<b>Mouse Swiss Prot No :</b>	Q8VDN2
<b>Rat Gene Id :</b>	24211
<b>Rat Swiss Prot No :</b>	P06685
<b>Immunogen :</b>	The antiserum was produced against synthesized peptide derived from human ATPase around the phosphorylation site of Ser16. AA range:5-54
<b>Specificity :</b>	Phospho-Na <sup>+</sup> /K <sup>+</sup> -ATPase α1 (S16) Polyclonal Antibody detects endogenous levels of Na <sup>+</sup> /K <sup>+</sup> -ATPase α1 protein only when phosphorylated at S16.


<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:5000.. IF 1:50-200
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Observed Band :</b>	112kD
<b>Cell Pathway :</b>	Cardiac muscle contraction;Aldosterone-regulated sodium reabsorption;
<b>Background :</b>	<p>The protein encoded by this gene belongs to the family of P-type cation transport ATPases, and to the subfamily of Na<sup>+</sup>/K<sup>+</sup> -ATPases. Na<sup>+</sup>/K<sup>+</sup> -ATPase is an integral membrane protein responsible for establishing and maintaining the electrochemical gradients of Na and K ions across the plasma membrane. These gradients are essential for osmoregulation, for sodium-coupled transport of a variety of organic and inorganic molecules, and for electrical excitability of nerve and muscle. This enzyme is composed of two subunits, a large catalytic subunit (alpha) and a smaller glycoprotein subunit (beta). The catalytic subunit of Na<sup>+</sup>/K<sup>+</sup> -ATPase is encoded by multiple genes. This gene encodes an alpha 1 subunit. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, May 2009],</p>
<b>Function :</b>	<p>catalytic activity:ATP + H(2)O + Na(+)(In) + K(+)(Out) = ADP + phosphate + Na(+)(Out) + K(+)(In).,function:This is the catalytic component of the active enzyme, which catalyzes the hydrolysis of ATP coupled with the exchange of sodium and potassium ions across the plasma membrane. This action creates the electrochemical gradient of sodium and potassium ions, providing the energy for active transport of various nutrients.,PTM:Phosphorylation on Tyr-10 modulates pumping activity.,similarity:Belongs to the cation transport ATPase (P-type) family.,similarity:Belongs to the cation transport ATPase (P-type) family. Type IIC subfamily.,subcellular location:Identified by mass spectrometry in melanosome fractions from stage I to stage IV.,subunit:Composed of three subunits: alpha (catalytic), beta and gamma. Binds the HLA class II histocompatibility antigen, DR1.,</p>
<b>Subcellular Location :</b>	<p>Basolateral cell membrane ; Multi-pass membrane protein . Cell membrane, sarcolemma ; Multi-pass membrane protein . Cell projection, axon . Melanosome . Identified by mass spectrometry in melanosome fractions from stage I to stage IV. .</p>

**Expression :**

Brain,Cerebellum,Cervix,Placenta,Retinal pigment epithelium

**Products Images**


Immunohistochemistry analysis of paraffin-embedded human lung carcinoma, using ATPase (Phospho-Ser16) Antibody. The picture on the right is blocked with the phospho peptide.

ATPase  
(pSer16)


Western blot analysis of lysates from 293 cells treated with PMA 125ng/ml 30', using ATPase (Phospho-Ser16) Antibody. The lane on the right is blocked with the phospho peptide.