

## EGF Receptor (Phospho Tyr845) rabbit pAb

Catalog No: YP1325

**Reactivity:** Human;Rat;Mouse;

**Applications:** WB

Target: EGFR

**Fields:** >>EGFR tyrosine kinase inhibitor resistance;>>Endocrine resistance;>>MAPK

signaling pathway;>>ErbB signaling pathway;>>Ras signaling pathway;>>Rap1

signaling pathway;>>Calcium signaling pathway;>>HIF-1 signaling pathway;>>FoxO signaling pathway;>>Phospholipase D signaling pathway;>>Endocytosis;>>PI3K-Akt signaling pathway;>>Focal adhesion;>>Adherens junction;>>Gap junction;>>JAK-STAT signaling

pathway;>>Regulation of actin cytoskeleton;>>GnRH signaling

pathway;>>Estrogen signaling pathway;>>Oxytocin signaling pathway;>>Relaxin

signaling pathway;>>Parathyroid hormone synthesis, secretion and

action;>>Cushing syndrome;>>Epithelial cell signaling in Helicobacter pylori

infection;>>Shigellosis;>>Hepatitis C;>>Human cytomegalovirus infection;>>Human papillomavirus infection;>>Coronavirus disease -

COVID-19;>>Pathways in cancer;>>Proteoglycans in cancer;>>MicroRNAs in

cancer;>>Chemical carcinogenesis - receptor activation;>>Chemical

carcinogenesis - reactive oxygen species;>>Colorectal cance

Gene Name: EGFR ERBB ERBB1 HER1

**Protein Name:** EGF Receptor (Tyr845)

Human Gene Id: 1956

**Human Swiss Prot** 

No:

Mouse Gene Id: 13649

**Mouse Swiss Prot** 

No:

Immunogen: Synthesized phosho peptide around human EGF Receptor (Tyr845)

**Specificity:** This antibody detects endogenous levels of Human EGF Receptor (phospho-

Tyr845)

P00533

Q01279



**Formulation:** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

**Dilution:** WB 1:1000-2000

**Purification:** The antibody was affinity-purified from rabbit serum by affinity-chromatography

using specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 134kD

Cell Pathway: MAPK\_ERK\_Growth;MAPK\_G\_Protein;ErbB\_HER;Calcium;Cytokine-cytokine

receptor interaction; Endocytosis; Dorso-ventral axis formation; Focal adhesion; Adherens\_Junction; Gap junction; Regulates Actin and Cytosk

**Background:** The protein encoded by this gene is a transmembrane glycoprotein that is a

member of the protein kinase superfamily. This protein is a receptor for members of the epidermal growth factor family. EGFR is a cell surface protein that binds to epidermal growth factor. Binding of the protein to a ligand induces receptor dimerization and tyrosine autophosphorylation and leads to cell proliferation. Mutations in this gene are associated with lung cancer. [provided by RefSeq, Jun

2016],

**Function:** catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine

phosphate., disease: Defects in EGFR are associated with lung cancer [MIM:211980]., function: Isoform 2/truncated isoform may act as an

antagonist., function: Receptor for EGF, but also for other members of the EGF family, as TGF-alpha, amphiregulin, betacellulin, heparin-binding EGF-like growth factor, GP30 and vaccinia virus growth factor. Is involved in the control of cell growth and differentiation. Phosphorylates MUC1 in breast cancer cells and

increases the interaction of MUC1 with C-SRC and CTNNB1/beta-

catenin.,miscellaneous:Binding of EGF to the receptor leads to dimerization, internalization of the EGF-receptor complex, induction of the tyrosine kinase activity, stimulation of cell DNA synthesis, and cell proliferation.,online

information:EGFR entry,PTM:Monoubiquitinated and polyubiquitinated upon EGF

stimu

Subcellular Location:

Cell membrane; Single-pass type I membrane protein. Endoplasmic reticulum membrane; Single-pass type I membrane protein. Golgi apparatus membrane; Single-pass type I membrane protein. Nucleus membrane; Single-pass type I membrane protein. Endosome . Endosome membrane. Nucleus . In response to EGF, translocated from the cell membrane to the nucleus via Golgi and ER (PubMed:20674546, PubMed:17909029). Endocytosed upon activation by ligand



(PubMed:2790960, PubMed:17182860, PubMed:27153536, PubMed:17909029). Colocalized with GPER1 in the nucleus of estrogen agonist-

induced cancer-associated fibroblasts (CAF) (PubMed:20551055). .; [Isoform 2]: Secreted.

**Expression :** Ubiquitously expressed. Isoform 2 is also expressed in ovarian cancers.

## **Products Images**