

## HER2 (Phospho Tyr1139) Polyclonal Antibody


| Catalog No :        | YP1240  |
|---------------------|---|
| Reactivity :        | Human;Mouse;Rat   |
| Applications :      | IHC;IF;WB   |
| Target :            | HER2  |
| Fields :            | >>EGFR tyrosine kinase inhibitor resistance;>>Endocrine<br>resistance;>>Platinum drug resistance;>>MAPK signaling pathway;>>ErbB<br>signaling pathway;>>Calcium signaling pathway;>>HIF-1 signaling<br>pathway;>>PI3K-Akt signaling pathway;>>Focal adhesion;>>Adherens<br>junction;>>Tight junction;>>Pathways in cancer;>>Proteoglycans in<br>cancer;>>MicroRNAs in cancer;>>Pancreatic cancer;>>Endometrial<br>cancer;>>Prostate cancer;>>Bladder cancer;>>Non-small cell lung<br>cancer;>>Breast cancer;>>Gastric cancer;>>Central carbon metabolism in<br>cancer |
| Gene Name :         | ERBB2 HER2 MLN19 NEU NGL  |
| Protein Name :      | HER2 (Phospho-Tyr1139)  |
| Human Gene Id :     | 2064  |
| Human Swiss Prot    | P04626  |
| No :<br>Immunogen : | Synthesized peptide derived from human HER2 (Phospho-Tyr1139)   |
| Specificity :       | This antibody detects endogenous phospho levels of HER2 (Phospho-Tyr1139) at Human:Y1139, Mouse:Y1140, Rat:Y1141  |
| Formulation :       | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| Source :            | Polyclonal, Rabbit,IgG  |
| Dilution :          | IHC 1:50-200, WB 1:500-2000. IF 1:50-200  |
| Purification :      | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.   |



| Best Tools for immunology Research |  |  |
|------------------------------------|--|--|
| <b>Concentration :</b>             | 1 mg/ml  |  |
| Storage Stability :                | -15°C to -25°C/1 year(Do not lower than -25°C)   |  |
| <b>Observed Band :</b>             | 180kD  |  |
| Background :                       | This gene encodes a member of the epidermal growth factor (EGF) receptor<br>family of receptor tyrosine kinases. This protein has no ligand binding domain of<br>its own and therefore cannot bind growth factors. However, it does bind tightly to<br>other ligand-bound EGF receptor family members to form a heterodimer,<br>stabilizing ligand binding and enhancing kinase-mediated activation of<br>downstream signalling pathways, such as those involving mitogen-activated<br>protein kinase and phosphatidylinositol-3 kinase. Allelic variations at amino acid<br>positions 654 and 655 of isoform a (positions 624 and 625 of isoform b) have<br>been reported, with the most common allele, lle654/lle655, shown here.<br>Amplification and/or overexpression of this gene has been reported in numerous<br>cancers, including breast and ovarian tumors. Alternative splicing results in<br>several additional transcript variants, some encoding d        |  |
| Function :                         | catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine<br>phosphate.,disease:Defects in ERBB2 are associated with familial glioma of brain<br>[MIM:137800]; also called glioblastoma multiforme. Gliomas are central nervous<br>system neoplasms derived from glial cells and comprise astrocytomas,<br>glioblastoma multiforme, oligodendrogliomas, and<br>ependymomas.,disease:Defects in ERBB2 are associated with gastric cancer<br>[MIM:137215]; also known as hereditary familial diffuse gastric cancer<br>(HDGC).,disease:Defects in ERBB2 are associated with lung cancer<br>[MIM:211980]; also called adenocarcinoma of lung.,disease:Defects in ERBB2<br>are associated with ovarian cancer [MIM:167000]. Ovarian cancer is the leading<br>cause of death from gynecologic malignancy. It is characterized by advanced<br>presentation with loco-regional dissemination in the peritoneal cavity and the rare<br>incidence of viscera |  |
| Subcellular<br>Location :          | [Isoform 1]: Cell membrane ; Single-pass type I membrane protein. Early<br>endosome . Cytoplasm, perinuclear region. Nucleus. Translocation to the nucleus<br>requires endocytosis, probably endosomal sorting and is mediated by importin<br>beta-1/KPNB1. Also detected in VPS35-positive endosome-to-TGN retrograde<br>vesicles (PubMed:31138794); [Isoform 2]: Cytoplasm. Nucleus.; [Isoform 3]:<br>Cytoplasm. Nucleus.  |  |
| Expression :                       | Expressed in a variety of tumor tissues including primary breast tumors and tumors from small bowel, esophagus, kidney and mouth.  |  |

## Products Images





Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).