


KIF1B (Phospho Ser1487) rabbit pAb

Catalog No :	YP1379
Reactivity :	Human;Rat;Mouse;
Applications :	WB;ELISA;IHC
Target :	KIF1B
Gene Name :	KIF1B KIAA0591 KIAA1448
Protein Name :	KIF1B (Ser1487)
Human Gene Id :	23095
Human Swiss Prot No :	O60333
Mouse Gene Id :	16561
Mouse Swiss Prot No :	Q60575
Rat Gene Id :	117548
Rat Swiss Prot No :	O88658
Immunogen :	Synthesized phospho peptide around human KIF1B (Ser1487)
Specificity :	This antibody detects endogenous levels of Human KIF1B (phospho-Ser1487)
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000;IHC 1:50-300; ELISA 2000-20000
Purification :	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.

Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	200kD
Background :	This gene encodes a motor protein that transports mitochondria and synaptic vesicle precursors. Mutations in this gene cause Charcot-Marie-Tooth disease, type 2A1. [provided by RefSeq, Jul 2008],
Function :	disease:Defects in KIF1B are the cause of Charcot-Marie-Tooth disease type 2A1 (CMT2A1) [MIM:118210]. CMT2A1 is a form of Charcot-Marie-Tooth disease, the most common inherited disorder of the peripheral nervous system. Charcot-Marie-Tooth disease is classified in two main groups on the basis of electrophysiologic properties and histopathology: primary peripheral demyelinating neuropathy or CMT1, and primary peripheral axonal neuropathy or CMT2. Neuropathies of the CMT2 group are characterized by signs of axonal regeneration in the absence of obvious myelin alterations, normal or slightly reduced nerve conduction velocities, and progressive distal muscle weakness and atrophy.,function:Motor for anterograde transport of mitochondria. Has a microtubule plus end-directed motility.,similarity:Belongs to the kinesin-like protein family.,similarity:Belongs to the kinesin-like protein family. U
Subcellular Location :	Cytoplasm, cytoskeleton. Mitochondrion . Cell projection, axon .; [Isoform 1]: Cytoplasmic vesicle, secretory vesicle, synaptic vesicle .
Expression :	Isoform 3 is abundant in the skeletal muscle. It is also expressed in fetal brain, lung and kidney, and adult heart, placenta, testis, ovary and small intestine. Isoform 2 is abundant in the brain and also expressed in fetal heart, lung, liver and kidney, and adult skeletal muscle, placenta, liver, kidney, heart, spleen, thymus, prostate, testis, ovary, small intestine, colon and pancreas.

Products Images



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).