

## Rac1/CDC42 (Phospho Ser71) rabbit pAb

Catalog No: YP1620

**Reactivity:** Human; Mouse; Rat

**Applications:** WB;ELISA;IHC

Target: Rac1/CDC42

**Fields:** >>MAPK signaling pathway;>>Ras signaling pathway;>>Rap1 signaling

pathway;>>cAMP signaling pathway;>>Chemokine signaling

pathway;>>Sphingolipid signaling pathway;>>Phagosome;>>PI3K-Akt signaling

pathway;>>Wnt signaling pathway;>>Axon guidance;>>VEGF signaling pathway;>>Osteoclast differentiation;>>Focal adhesion;>>Adherens

junction;>>Tight junction;>>Neutrophil extracellular trap formation;>>Toll-like receptor signaling pathway;>>Natural killer cell mediated cytotoxicity;>>B cell receptor signaling pathway;>>Fc epsilon RI signaling pathway;>>Fc gamma R-mediated phagocytosis;>>Leukocyte transendothelial migration;>>Neurotrophin signaling pathway;>>Regulation of actin cytoskeleton;>>Non-alcoholic fatty liver disease;>>AGE-RAGE signaling pathway in diabetic complications;>>Pancreatic

secretion;>>Amyotrophic lateral sclerosis;>>Prion disease;>>Pathways of neurodegeneration - multiple diseases;>>Bacterial invasion of epithelial

cells;>>Epithelial cell signaling in Helicobacter pylori infect

Gene Name: RAC1 TC25 MIG5

Protein Name: Rac1/CDC42 (Phospho Ser71)

Human Gene Id: 5879

**Human Swiss Prot** P63000/P15153/P60763/P60953

No:

Mouse Gene ld: 19353

Mouse Swiss Prot P63001

No:

**Rat Gene Id :** \_\_\_363875

Rat Swiss Prot No : Q6RUV5



Immunogen: Synthesized peptide derived from human Rac1/CDC42 (Phospho Ser71)

**Specificity:** This antibody detects endogenous levels of Human, Mouse, Rat Rac1/CDC42

(Phospho Ser71)

**Formulation :** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

**Dilution :** WB 1:500-2000;IHC 1:50-300; ELISA 2000-20000

**Purification:** The antibody was affinity-purified from rabbit serum by affinity-chromatography

using specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight: 21kD

Subcellular Location:

Cell membrane; Lipid-anchor; Cytoplasmic side. Melanosome. Cytoplasm. Cell projection, lamellipodium. Cell projection, dendrite. Cell junction, synapse. Nucleus. Inner surface of plasma membrane possibly with attachment requiring prenylation of the C-terminal cysteine (PubMed:1903399). Identified by mass

spectrometry in melanosome fractions from stage I to stage IV

(PubMed:17081065). Found in the ruffled border (a late endosomal-like

compartment in the plasma membrane) of bone-resorbing osteoclasts. Localizes

to the lamellipodium in a SH3RF1-dependent manner (By similarity). In macrophages, cytoplasmic location increases upon CSF1 stimulation (By similarity). Activation by GTP-binding promotes nuclear localization

(PubMed:12551911)...


**Expression:** Isoform B is predominantly identified in skin and epithelial tissues from the

intestinal tract. Its expression is elevated in colorectal tumors at various stages of


neoplastic progression, as compared to their respective adjacent tissues.

## **Products Images**


2/4




Western blot analysis of Rac1(Phospho-Ser71) antibody in 293 cells lysates, cell treated or untreated with PMA 125ng/ml 30', 4° over night, secondary antibody(cat: RS0002 was diluted at 1:10000, 37° 1hour.



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using Rac1(Phospho-Ser71) antibody



Immunohistochemical analysis of paraffin-embedded human uterus. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).