

CD142 (PN0276) Nb-FC recombinant antibody

Catalog No: YA0102

Reactivity: Human

Applications: ELISA

Target: CD142

Gene Name: F3

Protein Name : Tissue factor (TF) (Coagulation factor III) (Thromboplastin) (CD antigen CD142)

Human Gene ld: 2152

Human Swiss Prot

No:

Immunogen: Purified recombinant Human CD142

P13726

Specificity: This recombinant monoclonal antibody can detects endogenous levels of CD142

protein.

Formulation: Phosphate-buffered solution

Source: Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain,

recombinantly produced from 293F cell

Dilution: ELISA 1:5000-100000

Purification: Recombinant Expression and Affinity purified

Concentration: Please check the information on the tube

Storage Stability: -15°C to -25°C/1 year(Avoid freeze / thaw cycles)

Background: CD142, also known as Tissue Factor (TF), Coagulation Factor III, and

Thromboplastin, is a 45 kD type I transmembrane glycoprotein. It is expressed on the surface of a variety of cells that are physically separated from the circulating blood which include smooth muscle cells, fibroblasts, keratinocytes, glomerular

epithelial cells (cytoplasmic inclusions), astrocytes, myocardium, liver stromal cells, pancreas cells, and is also expressed on activated monocytes and stimulated endothelial cells. CD142 is a high-affinity receptor for coagulation factor VII and initiates the extrinsic pathway of blood coagulation. CD142 also plays an important role in a variety of diseases such as sepsis, atherosclerosis, and cancer.

Function:

Initiates blood coagulation by forming a complex with circulating factor VII or VIIa. The [TF:VIIa] complex activates factors IX or X by specific limited proteolysis. TF plays a role in normal hemostasis by initiating the cell-surface assembly and propagation of the coagulation protease cascade.

Subcellular Location:

[Isoform 1]: Membrane ; Single-pass type I membrane protein .; [Isoform 2]: Secreted .

Expression: Lung, placenta and pancreas.

Products Images

