

CD166 (PN0357) Nb-FC recombinant antibody

Catalog No :	YA0131
Reactivity :	Human
Applications :	ELISA
Target :	CD166
Gene Name :	ALCAM MEMD
Protein Name :	CD166 antigen (Activated leukocyte cell adhesion molecule) (CD antigen CD166)
Human Gene Id :	214
Human Swiss Prot No :	Q13740
Immunogen :	Purified recombinant Human CD166
Specificity :	This recombinant monoclonal antibody can detects endogenous levels of CD166 protein.
Formulation :	Phosphate-buffered solution
Source :	Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain , recombinantly produced from 293F cell
Dilution :	ELISA 1:5000-100000
Purification :	Recombinant Expression and Affinity purified
Concentration :	Please check the information on the tube
Storage Stability :	-15°C to -25°C/1 year(Avoid freeze / thaw cycles)
Cell Pathway :	Cell adhesion molecules (CAMs);
Background :	This gene encodes activated leukocyte cell adhesion molecule (ALCAM), also

known as CD166 (cluster of differentiation 166), which is a member of a subfamily of immunoglobulin receptors with five immunoglobulin-like domains (VVC2C2C2) in the extracellular domain. This protein binds to T-cell differentiation antigen CD6, and is implicated in the processes of cell adhesion and migration. Multiple alternatively spliced transcript variants encoding different isoforms have been found. [provided by RefSeq, Aug 2011]

Function :

domain: The CD6 binding site is located in the N-terminal Ig-like domain. Cell adhesion molecule that binds to CD6. Involved in neurite extension by neurons via heterophilic and homophilic interactions. May play a role in the binding of T- and B-cells to activated leukocytes, as well as in interactions between cells of the nervous system. similarity: Contains 2 Ig-like V-type (immunoglobulin-like) domains. similarity: Contains 3 Ig-like C2-type (immunoglobulin-like) domains. tissue specificity: Spleen, placenta, liver, and weakly in liver. Expressed by activated T-cells, B-cells, monocytes and thymic epithelial cells. Expressed by neurons in the brain. Restricted expression in tumor cell lines. Preferentially expressed in highly metastasizing melanoma cell lines.

Subcellular Location :

Cell membrane ; Single-pass type I membrane protein . Cell projection, axon . Cell projection, dendrite . Detected at the immunological synapse, i.e, at the contact zone between antigen-presenting dendritic cells and T-cells (PubMed:15294938, PubMed:16352806). Colocalizes with CD6 and the TCR/CD3 complex at the immunological synapse (PubMed:15294938). .; [Isoform 3]: Secreted .

Expression :

Detected on hematopoietic stem cells derived from umbilical cord blood (PubMed:2474813). Detected on lymph vessel endothelial cells, skin and tonsil (PubMed:23169771). Detected on peripheral blood monocytes (PubMed:154873). Detected on monocyte-derived dendritic cells (at protein level) (PubMed:1635286). Detected at low levels in spleen, placenta, liver (PubMed:952422). Expressed by activated T-cells, B-cells, monocytes and thymic epithelial cells (PubMed:7767). Isoform 1 and isoform 3 are detected in vein and artery endothelial cells, astrocytes, keratinocytes and artery smooth muscle cells (PubMed:15496415). Expressed by neurons in the brain. Restricted expression in tumor cell lines. Detected in highly metastasizing melanoma cell lines (PubMed:952422).

Products Images

