

CD71 (PN0514) Nb-FC recombinant antibody

Catalog No :	YA0481
Reactivity :	Human
Applications :	ELISA
Target :	CD71
Gene Name :	TFRC
Protein Name :	Transferrin receptor protein 1 (TR) (TfR) (TfR1) (Trfr) (T9) (p90) (CD antigen CD71) [Cleaved into: Transferrin receptor protein 1, serum form (sTfR)]
Human Gene Id :	7037
Human Swiss Prot No :	P02786
Immunogen :	Purified recombinant Human CD71
Specificity :	This recombinant monoclonal antibody can detects endogenous levels of CD71 protein.
Formulation :	Phosphate-buffered solution
Source :	Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain , recombinantly produced from 293F cell
Dilution :	ELISA 1:5000-100000
Purification :	Recombinant Expression and Affinity purified
Concentration :	Please check the information on the tube
Storage Stability :	-15°C to -25°C/1 year(Avoid freeze / thaw cycles)
Background :	transferrin receptor(TFRC) Homo sapiens This gene encodes a cell surface receptor necessary for cellular iron uptake by the process of receptor-mediated endocytosis.This receptor is required for erythropoiesis and neurologic

development. Multiple alternatively spliced variants have been identified.
[provided by RefSeq, Sep 2015]

Function :

Cellular uptake of iron occurs via receptor-mediated endocytosis of ligand-occupied transferrin receptor into specialized endosomes . Endosomal acidification leads to iron release. The apotransferrin-receptor complex is then recycled to the cell surface with a return to neutral pH and the concomitant loss of affinity of apotransferrin for its receptor. Transferrin receptor is necessary for development of erythrocytes and the nervous system (By similarity). A second ligand, the heditary hemochromatosis protein HFE, competes for binding with transferrin for an overlapping C-terminal binding site. Positively regulates T and B cell proliferation through iron uptake . Acts as a lipid sensor that regulates mitochondrial fusion by regulating activation of the JNK pathway . When dietary levels of stearate (C18:0) are low, promotes activation of the JNK pathway, resulting in HUWE1-mediated ubiqu

Subcellular Location :

Cell membrane ; Single-pass type II membrane protein . Melanosome . Identified by mass spectrometry in melanosome fractions from stage I to stage IV. .; [Transferrin receptor protein 1, serum form]: Secreted .

Products Images