

MAP2(N-term) mouse mAb

Catalog No: YM1513

Reactivity: Human; Mouse

Applications: WB;IP;ICC

Target: MAP-2

Gene Name: map2

Human Gene Id: 4133

Human Swiss Prot

No:

Mouse Swiss Prot

No:

Immunogen: Purified recombinant human MAP2 (N-terminus) protein fragments expressed in

E.coli

P11137

P20357

Specificity: This antibody detects endogenous levels of MAP2 (N-terminus) and does not

cross-react with related proteins.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Monoclonal, Mouse

Dilution: wb dilution 1:1000 icc dilution 1:150

Purification: The antibody was affinity-purified from mouse ascites by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 300kD

Background: This gene encodes a protein that belongs to the microtubule-associated protein

1/3



family. The proteins of this family are thought to be involved in microtubule assembly, which is an essential step in neurogenesis. The products of similar genes in rat and mouse are neuron-specific cytoskeletal proteins that are enriched in dentrites, implicating a role in determining and stabilizing dentritic shape during neuron development. A number of alternatively spliced variants encoding distinct isoforms have been described. [provided by RefSeq, Jan 2010],

Function:

alternative products:Additional isoforms seem to exist,function:The exact function of MAP2 is unknown but MAPs may stabilize the microtubules against depolymerization. They also seem to have a stiffening effect on microtubules.,PTM:MAP2A/c is phosphorylated. Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 3 Tau/MAP repeats.,similarity:Contains 4 Tau/MAP repeats.,


Subcellular Location :

Cytoplasm, cytoskeleton . Cell projection, dendrite .


Expression:

Brain, Brain cortex, Epithelium, Pancreas, Testis,


Products Images



Western blot detection of MAP2(N-terminus) in Mouse Brain tissue and Hela cell lysates using MAP2(N-terminus) mouse mAb (1:1000 diluted). Predicted band size: 202KDa. Observed band size: 300KDa.



Immunocytochemistry staining of Hela cells fixed with 4% Paraformaldehyde and using anti-MAP2(N-term) mouse mAb (dilution 1:100).



Immunoprecipitation analysis of Hela cell lysates using MAP2 (Nterminus) mouse mAb.