

## FITC\* Goat Anti Rabbit IgG(H+L)

Catalog No: RS0004

Reactivity: Rabbit

**Applications:** IF;ELISA

**Target:** Rabbit IgG(H+L)

**Formulation :** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Goat

**Dilution:** Optimal working dilutions should be determined experimentally by the

investigator. Suggested starting dilutions are IF (1:250 - 1:2000),

**Purification:** The antibody was affinity-purified from goat antiserum by affinity-

chromatography using epitope-specific immunogen.

**Concentration:** 4 mg/ml

**Storage Stability:** -15°C to -25°C/1 year(Do not lower than -25°C)


**Background:** Immunoway secondary antibodies are available conjugated to enzyme, biotin or

fluorophore for use in a variety of antibody-based applications including Western Blot, ImmunoHistoChemistry, ImmunoFluorescence, Flow Cytometry and ELISA. We offer high quality secondary antibodies from goat, rabbit and donkey sources

for your each application. Serum adsorbed secondary antibodies are also available and are recommended for use with immunoglobulin-rich samples.

## **Products Images**

1/2



Zhang, Zi-Bin, et al. "miRNA-7a-2-3p Inhibits Neuronal Apoptosis in Oxygen-Glucose Deprivation (OGD) Model." Frontiers in neuroscience 13 (2019).