

## **SOCS4 Polyclonal Antibody**

Catalog No: YN1373

**Reactivity:** Human; Mouse

**Applications:** WB;ELISA

Target: SOCS4

**Fields:** >>JAK-STAT signaling pathway;>>Insulin signaling pathway;>>Prolactin

signaling pathway;>>Type II diabetes mellitus

Gene Name: SOCS4 SOCS7

**Protein Name:** Suppressor of cytokine signaling 4 (SOCS-4) (Suppressor of cytokine signaling

7) (SOCS-7)

Q91ZA6

Human Gene Id: 122809

Human Swiss Prot Q8WXH5

No:

**Mouse Swiss Prot** 

No:

**Immunogen:** Synthesized peptide derived from part region of human protein

**Specificity:** SOCS4 Polyclonal Antibody detects endogenous levels of protein.

**Formulation :** Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

**Dilution:** WB 1:500-2000 ELISA 1:5000-20000

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

**Storage Stability:** -15°C to -25°C/1 year(Do not lower than -25°C)

1/2



**Observed Band:** 48kD

**Cell Pathway:** Jak\_STAT;Insulin\_Receptor;Type II diabetes mellitus;

**Background:** The protein encoded by this gene contains a SH2 domain and a SOCS BOX

domain. The protein thus belongs to the suppressor of cytokine signaling (SOCS), also known as STAT-induced STAT inhibitor (SSI), protein family. SOCS family members are known to be cytokine-inducible negative regulators of cytokine signaling. Two alternatively spliced transcript variants encoding the same protein

have been found for this gene. [provided by RefSeq, Jul 2008],

**Function:** domain: The SOCS box domain mediates the interaction with the Elongin BC

complex, an adapter module in different E3 ubiquitin ligase

complexes.,function:SOCS family proteins form part of a classical negative feedback system that regulates cytokine signal transduction. May be a substrate recognition component of a SCF-like ECS (Elongin BC-CUL2/5-SOCS-box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and

subsequent proteasomal degradation of target proteins..pathway:Protein

modification; protein ubiquitination., similarity: Contains 1 SH2

domain., similarity: Contains 1 SOCS box domain.,

Subcellular Location:

intracellular, cytoplasm,

**Expression:** Endometrium carcinoma cell line, Neuroblastoma, Placenta,

## **Products Images**

2/2