

SOCS5 Polyclonal Antibody

Catalog No :	YN1374
Reactivity :	Human;Mouse
Applications :	WB;ELISA
Target :	SOCS5
Fields :	>>JAK-STAT signaling pathway;>>Prolactin signaling pathway
Gene Name :	SOCS5 CIS6 CISH5 CISH6 KIAA0671
Protein Name :	Suppressor of cytokine signaling 5 (SOCS-5) (Cytokine-inducible SH2 protein 6) (CIS-6) (Cytokine-inducible SH2-containing protein 5)
Human Gene Id :	9655
Human Swiss Prot No :	O75159
Mouse Swiss Prot No :	O54928
Immunogen :	Synthesized peptide derived from part region of human protein
Specificity :	SOCS5 Polyclonal Antibody detects endogenous levels of protein.
Formulation :	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000 ELISA 1:5000-20000
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band :	58kD
Cell Pathway :	Jak_STAT;
Background :	<p>The protein encoded by this gene contains a SH2 domain and a SOCS BOX domain. The protein thus belongs to the suppressor of cytokine signaling (SOCS) family, also known as STAT-induced STAT inhibitor (SSI) protein family. SOCS family members are known to be cytokine-inducible negative regulators of cytokine signaling. The specific function of this protein has not yet been determined. Two alternatively spliced transcript variants encoding an identical protein have been reported. [provided by RefSeq, Jul 2008],</p>
Function :	<p>domain:The SOCS box domain mediates the interaction with the Elongin BC complex, an adapter module in different E3 ubiquitin ligase complexes.,function:SOCS family proteins form part of a classical negative feedback system that regulates cytokine signal transduction. SOCS5 is involved in regulating T-helper cell differentiation by inhibition of the IL4 signaling pathway which promotes differentiation into the Th2 phenotype. Can also partially inhibit IL6 and LIF signaling (By similarity). May be a substrate-recognition component of a SCF-like ECS (Elongin BC-CUL2/5-SOCS-box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins.,pathway:Protein modification; protein ubiquitination.,similarity:Contains 1 SH2 domain.,similarity:Contains 1 SOCS box domain.,subunit:Interacts, via the first 50 AA, with IL4RA (th</p>
Subcellular Location :	cytoplasm,
Expression :	Brain,Placenta,Testis,Thalamus,

Products Images