

TBB1 Polyclonal Antibody

Catalog No: YN1906

Reactivity: Human; Mouse

Applications: WB;ELISA

Target: Tubulin β

Fields: >>Phagosome;>>Gap junction;>>Alzheimer disease;>>Parkinson

disease;>>Amyotrophic lateral sclerosis;>>Huntington disease;>>Prion disease;>>Pathways of neurodegeneration - multiple diseases;>>Pathogenic

Escherichia coli infection;>>Salmonella infection

Gene Name: TUBB1

Protein Name: Tubulin beta-1 chain

Human Gene Id: 81027

Human Swiss Prot

·

No:

Mouse Swiss Prot

No:

Immunogen:

Synthesized peptide derived from part region of human protein AA range:

257-307

Q9H4B7

A2AQ07

Specificity: TBB1 Polyclonal Antibody detects endogenous levels of protein.

Formulation: Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1:500-2000 ELISA 1:5000-20000

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml



Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 49kD

Cell Pathway: Gap junction; Pathogenic Escherichia coli infection;

Background: This gene encodes a member of the beta tubulin protein family. Beta tubulins are

one of two core protein families (alpha and beta tubulins) that heterodimerize and assemble to form microtubules. This protein is specifically expressed in platelets and megakaryocytes and may be involved in proplatelet production and platelet release. A mutations in this gene is associated with autosomal dominant

macrothrombocytopenia. Two pseudogenes of this gene are found on

chromosome Y.[provided by RefSeq, Jul 2010],

Function: function: Tubulin is the major constituent of microtubules. It binds two moles of

GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha-chain.,similarity:Belongs to the tubulin

family., subunit: Dimer of alpha and beta chains.,

Subcellular Location:

Cytoplasm, cytoskeleton.

Expression: Hematopoietic cell-specific. Major isotype in leukocytes, where it represents

50% of all beta-tubulins.

Products Images

2/2